

Fuzzifying Allen’s Temporal Interval Relations

Steven Schockaert, Martine De Cock, and Etienne E. Kerre

Abstract—When the time span of an event is imprecise, it can be represented by a fuzzy set, called a fuzzy time interval. In this paper we propose a framework to represent, compute and reason about temporal relationships between such events. Since our model is based on fuzzy orderings of time points, it is not only suitable to express precise relationships between imprecise events (“Roosevelt died *before* the beginning of the Cold War”) but also imprecise relationships (“Roosevelt died *just before* the beginning of the Cold War”). We show that, unlike previous models, our model is a generalization that preserves many of the properties of the 13 relations Allen introduced for crisp time intervals. Furthermore, we show how our model can be used for efficient fuzzy temporal reasoning by means of a transitivity table. Finally, we illustrate its use in the context of question answering systems.

Index Terms—Temporal Reasoning, Fuzzy Relation, Fuzzy Ordering, Interval Algebra, Question Answering

I. INTRODUCTION

TEMPORAL representation and reasoning is an important facet in the design of many intelligent systems. For example, question answering systems require at least some basic temporal representation scheme to answer simple temporal questions such as “When was Franklin Roosevelt born?”. To enable question answering systems to answer more complex temporal questions, considerable effort has been done to extract temporal information from natural language texts (e.g., [1], [12], [15], [16], [23], [24], [25]) and to analyse complex temporal questions (e.g., [22]). However, temporal relationships expressed in natural language are often vague, e.g., “Roosevelt died *just before* the end of the Second World War.”. Moreover, historic time periods are more often than not characterized by a gradual beginning and/or ending [17]. The traditional temporal reasoning formalisms need to be extended to cope with this kind of vagueness which is inherently associated with real-world temporal information.

One of those well known formalisms is Allen’s temporal interval algebra [3]. Allen defined a set of 13 qualitative relations that may hold between two compact intervals $A = [a^-, a^+]$ and $B = [b^-, b^+]$. Table I shows how Allen expressed these precise relations by means of constraints on the boundaries of the crisp intervals involved. In this paper we extend Allen’s work to a more general formalism that can handle precise as well as imprecise relationships between crisp as well as fuzzy intervals.

Our first concern is generalizing the definitions of the qualitative relations of Table I to make them applicable to fuzzy intervals as opposed to only crisp intervals. Indeed, when

Steven Schockaert*, Martine De Cock and Etienne E. Kerre are with the Department of Applied Mathematics and Computer Science, Ghent University, Krijgslaan 281 (S9), B-9000 Gent, Belgium (e-mail: {Steven.Schockaert, Martine.DeCock, Etienne.Kerre}@UGent.be.)

*Research assistant of the Research Foundation – Flanders

TABLE I

ALLEN’S TEMPORAL INTERVAL RELATIONS BETWEEN INTERVALS
 $A = [a^-, a^+]$ AND $B = [b^-, b^+]$.

Name	Definition
before	$b(A, B) \equiv a^+ < b^-$
overlaps	$o(A, B) \equiv a^- < b^-$ and $b^- < a^+$ and $a^+ < b^+$
during	$d(A, B) \equiv b^- < a^-$ and $a^+ < b^+$
meets	$m(A, B) \equiv a^+ = b^-$
starts	$s(A, B) \equiv a^- = b^-$ and $a^+ < b^+$
finishes	$f(A, B) \equiv a^+ = b^+$ and $b^- < a^-$
equals	$e(A, B) \equiv a^- = b^-$ and $a^+ = b^+$
after	$bi(A, B) \equiv b(B, A)$
overlapped-by	$oi(A, B) \equiv o(B, A)$
contains	$di(A, B) \equiv d(B, A)$
met-by	$mi(A, B) \equiv m(B, A)$
started-by	$si(A, B) \equiv s(B, A)$
finished-by	$fi(A, B) \equiv f(B, A)$

an event is characterized by a gradual beginning and/or ending, it is natural to represent the corresponding time span as a fuzzy set, which we call a fuzzy (time) interval. Depending on the intended application, this fuzzy set can either be defined by an expert (e.g., [17], [20]), or it can be constructed automatically (e.g., [27]). Since we can not refer to the gradual beginning and endings of a fuzzy interval in the same way we refer to the boundaries of a crisp interval, we first have to provide a way to express that, for instance, the beginning of a fuzzy interval A is before the beginning of a fuzzy interval B (as needed in the definition of the qualitative relation “overlaps”). We suggest to do this by measuring the highest extent to which there exist a time point in A that occurs before all the time points in B . In general, in our approach, qualitative relations between fuzzy intervals are defined in terms of the ordering of the gradual beginning and endings of these intervals, which in turn are defined in terms of the ordering of the time points belonging to these intervals. The resulting qualitative relations between the fuzzy intervals are gradual, i.e., they may hold to some degree only, hence the name fuzzy temporal interval relations. When A and B are crisp, our approach reduces to Allen’s work.

Our second goal is providing a means to model *imprecise* relations to be able to express that, for instance, event A took place *just before* event B , or that A occurred *long after* B . Although these kind of relations are not considered in Allen’s original model, in our approach we arrive at them quite elegantly by using imprecise orderings of time points in the model sketched above. The resulting approach is applicable again both to crisp and fuzzy time intervals.

The paper is organized as follows: in the next section we review related work concerning fuzzifications of Allen’s interval relations. In Section III, we show how imprecise relationships between time points can be modelled by using fuzzy

orderings. In Section IV we rely on relatedness measures for fuzzy sets to lift these imprecise orderings of time points into relationships between fuzzy time intervals [29]. This results in a generalization of Allen’s 13 interval relations that are also applicable when the time intervals are fuzzy. Furthermore, this framework is powerful enough to additionally express imprecise relationships that are not considered in Allen’s original model. We show that our model preserves many important properties regarding (ir)reflexivity, (a)symmetry and transitivity, and that our generalized definitions remain mutual exclusive and exhaustive. Moreover, in Section V, we discuss fuzzy temporal reasoning and introduce a transitivity table to derive new temporal knowledge in an efficient way. This transitivity table is a generalization of the transitivity table that was introduced by Freksa in [13], which shows that no transitivity properties are lost in our generalized framework. Section VI illustrates the usefulness of our approach within the context of question answering systems. Sections III and IV contain many new results that require a mathematical proof. To preserve the continuity of the main text, we present these proofs in appendices.

II. RELATED WORK

Most fuzzifications of Allen’s interval algebra deal with uncertainty rather than imprecision (e.g., [4], [9], [10], [11], [14]). These approaches assume that — in the face of complete knowledge — the time span of an event can always be modelled as a crisp (time) interval. For example, Dubois and Prade [9] represent a time interval as a pair of possibility distributions which define the possible values of the endpoints of the crisp interval. Using possibility theory, the possibility and necessity of each of the interval relations can then be calculated. This approach also allows to model imprecise relations such as “*A* was *long* before *B*”. In a different approach adopted by Dutta [11], time intervals are abstract entities and the possibility, for each interval i and each event e , that e occurs in i is defined. In [4] uncertainty regarding the temporal relations that hold between crisp time intervals is considered in order to reason with statements such as “the possibility that $m(A, B)$ holds is 0.6”. Guesgen et al. [14] proposed a similar approach based on the notion of a conceptual neighborhood, a notion originally introduced in [13].

Temporal information is expressed w.r.t. a certain level of granularity (e.g., years, days, seconds,...) which partitions the timeline. In [7] it is argued that the time span of events lies often skew to the cells of this partitioning. Therefore, a rough set approach is adopted in which the time span of an event is represented by a lower approximation consisting of the cells of the partitioning which are fully included in this time span, and an upper approximation consisting of the cells of the partitioning which at least partially overlap with this time span. The temporal interval relations are redefined, using a directed variant of the RCC calculus [21], to cope with these “rough time intervals”.

In [6] it is suggested to represent time intervals as fuzzy sets, but no definitions of the interval relations are given. Most relevant to our approach are definitely the work of Nagypál

and Motik [17] and the work of Ohlbach [20] which are concerned with generalizing Allen’s interval relations when the time span of an event is represented as a fuzzy set. However, these approaches suffer from a number of important disadvantages. For example, the relation “equals” defined in [17] is not reflexive in general; for a continuous fuzzy set A in \mathbb{R} , $e(A, A) = s(A, A) = f(A, A) = d(A, A) = 0.5$, while, taking into account Allen’s intended meaning of these relations, one would expect $e(A, A) = 1$ and $s(A, A) = f(A, A) = d(A, A) = 0$. Moreover, imprecise temporal relations cannot be expressed. An approach similar to [17] was suggested in [8] within the context of ranking fuzzy numbers. Ohlbach [20] suggests an alternative approach which allows to model imprecise temporal relations such as “*A* *more or less* finishes *B*” based on measures of overlap for fuzzy sets. However, this approach cannot handle imprecise temporal relations such as “*A* was *long* before *B*”. Moreover, as pointed out in [20], many of the (ir)reflexivity, (a)symmetry and transitivity properties of the original temporal relations are lost in this approach; hence it is not suitable for temporal reasoning. Imprecise temporal relations are also considered in [10]; however, only crisp intervals are considered in this approach.

III. FUZZY ORDERING OF TIME POINTS

A. Definitions

The fuzzy temporal interval relations that we will define in the next section are based on orderings between the time points contained in the intervals. Throughout this paper, we represent time points as real numbers. A real number can, for example, be interpreted as the number of milliseconds since January 1, 1970, or the number of years since 1900. Because we want to model imprecise temporal relations, we need a way to express for two time points a and b that a is *long* before b , that a is *just* before b , and that a and b occur at *approximately* the same time.

Let $\alpha \in \mathbb{R}$ and $\beta \in [0, +\infty[$, then the extent to which a is long before b (w.r.t. (α, β)) can be expressed by the fuzzy relation $L_{(\alpha, \beta)}^{\ll}$ in \mathbb{R} defined as [9]

$$L_{(\alpha, \beta)}^{\ll}(a, b) = \begin{cases} 1 & \text{if } b - a > \alpha + \beta \\ 0 & \text{if } b - a \leq \alpha \\ \frac{b - a - \alpha}{\beta} & \text{otherwise} \end{cases} \quad (1)$$

for all a and b in \mathbb{R} . The partial mapping $L_{(\alpha, \beta)}^{\ll}(\cdot, b)$ is depicted in Figure 1(a). The parameters α and β define how the concept ‘long before’ should be interpreted. For a time point to be long before b to degree 1, the time gap with b should be at least $\alpha + \beta$. If the time gap with b is smaller than α , the time point is long before b to degree 0. In between there is a gradual transition. Although it seems natural to impose that α is positive, for technical reasons we only require $\alpha \in \mathbb{R}$. Moreover, as pointed out by Ohlbach [20], in some applications it may be desirable for some $a > b$ to express that a is (long) before b to a (small) strictly positive degree, which is only possible in our approach if we allow negative α ’s. $L_{(\alpha, \beta)}^{\ll}$ is a generalization of the crisp strict ordering relation




Fig. 1. Fuzzy ordering of time points

$<$. Indeed, imposing $\alpha = \beta = 0$, we obtain $L_{(0,0)}^{\ll}(a, b) = 1$ if $a < b$ and $L_{(0,0)}^{\ll}(a, b) = 0$ otherwise.

The fuzzy relation $L_{(\alpha,\beta)}^{\lessgtr}$ in \mathbb{R} is defined as [9]

$$L_{(\alpha,\beta)}^{\lessgtr}(a, b) = 1 - L_{(\alpha,\beta)}^{\ll}(b, a) \quad (2)$$

for all a and b in \mathbb{R} . $L_{(\alpha,\beta)}^{\lessgtr}(a, b)$ represents the extent to which b is not ‘long before’ a (w.r.t. (α, β)), in other words, the extent to which a is before or at approximately the same time as b . It holds that

$$L_{(\alpha,\beta)}^{\lessgtr}(a, b) = \begin{cases} 1 & \text{if } b - a \geq -\alpha \\ 0 & \text{if } b - a < -\alpha - \beta \\ \frac{b - a + \alpha + \beta}{\beta} & \text{otherwise} \end{cases} \quad (3)$$

Moreover, $L_{(0,0)}^{\lessgtr}(a, b) = 1$ if $a \leq b$ and $L_{(0,0)}^{\lessgtr}(a, b) = 0$ otherwise, i.e. $L_{(\alpha,\beta)}^{\lessgtr}$ is a generalization of the crisp ordering \leq .

As will become clear in Section IV, we only need the fuzzy relations $L_{(\alpha,\beta)}^{\ll}$ and $L_{(\alpha,\beta)}^{\lessgtr}$ to model imprecise temporal interval relations. The degree $E_{(\alpha,\beta)}(a, b)$ to which a occurs at approximately the same time as b , and the degree $J_{(\alpha,\beta)}(a, b)$ to which a is just before b can easily be expressed using $L_{(\alpha,\beta)}^{\ll}$ and $L_{(\alpha,\beta)}^{\lessgtr}$, i.e.

$$E_{(\alpha,\beta)}(a, b) = \min(L_{(\alpha,\beta)}^{\lessgtr}(a, b), L_{(\alpha,\beta)}^{\lessgtr}(b, a)) \quad (4)$$

$$J_{(\alpha,\beta)}(a, b) = \min(E_{(\alpha,\beta)}(a, b), L_{(0,0)}^{\ll}(a, b)) \quad (5)$$

An overview of the 4 fuzzy relations between time points is given in Figure 1.

Example 1: Assume that a time point corresponds to the number of years since January 1900. Using $\alpha = 2$ and $\beta = 8$, we obtain for example:

$$L_{(2,8)}^{\ll}(20, 23) = 0.125 \quad E_{(2,8)}(20, 23) = 0.875$$

$$L_{(2,8)}^{\lessgtr}(20, 23) = 1 \quad J_{(2,8)}(20, 23) = 0.875$$

expressing that 20 occurred long before 23 to a low degree, that 20 occurred just before 23 to a high degree etc. On the other hand we also have:

$$L_{(2,8)}^{\ll}(23, 20) = 0 \quad E_{(2,8)}(23, 20) = 0.875$$

$$L_{(2,8)}^{\lessgtr}(23, 20) = 0.875 \quad J_{(2,8)}(23, 20) = 0$$

In other words, although 23 is not considered to be long before or just before 20, it is still considered to be before or at approximately the same time as 20 to a high degree.

B. Properties

The fuzzy relations $L_{(\alpha,\beta)}^{\ll}$ and $L_{(\alpha,\beta)}^{\lessgtr}$ behave as can be intuitively expected from orderings. First recall that in general a fuzzy relation R in a universe U is called

(1) reflexive iff $R(u, u) = 1$ for all u in U

(2) irreflexive iff $R(u, u) = 0$ for all u in U

(3) symmetric iff $R(u, v) = R(v, u)$ for all u and v in U

(4) T -asymmetric iff $T(R(u, v), R(v, u)) = 0$ for all u and v in U

(5) T -transitive iff $T(R(u, v), R(v, w)) \leq R(u, w)$ for all u, v and w in U

for an arbitrary triangular norm T . Furthermore, if a fuzzy relation E in U is reflexive, symmetric and T -transitive, E is called a fuzzy T -equivalence relation. A fuzzy T - E -ordering relation [8] is then defined as a T -transitive fuzzy relation L in U which is

- (6) E -reflexive i.e. $E(u, v) \leq L(u, v)$
for all u and v in U
- (7) T - E -antisymmetric i.e. $T(L(u, v), L(v, u)) \leq E(u, v)$
for all u and v in U

Transitivity of the fuzzy orderings is of particular importance for temporal reasoning. Many interesting properties regarding the transitivity of $L_{(\alpha, \beta)}^{\ll}$ and $L_{(\alpha, \beta)}^{\lessgtr}$ follow from an important characterization of their composition. Recall that in general the sup- T composition of two fuzzy relations R and S in U is the fuzzy relation $R \circ_T S$ in U defined for each u and w in U by

$$(R \circ_T S)(u, w) = \sup_{v \in U} T(R(u, v), S(v, w)) \quad (6)$$

Throughout this paper, we use T_W to denote the Łukasiewicz t-norm ,i.e.

$$T_W(x, y) = \max(0, x + y - 1)$$

for all x and y in $[0, 1]$.

Proposition 1 (Composition): Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $\beta_1, \beta_2 \geq 0$; it holds that

$$L_{\gamma_1}^{\ll} \circ_{T_W} L_{\gamma_2}^{\ll} = L_{(\alpha_1 + \alpha_2 + \min(\beta_1, \beta_2), \max(\beta_1, \beta_2))}^{\ll} \quad (7)$$

$$L_{\gamma_1}^{\lessgtr} \circ_{T_W} L_{\gamma_2}^{\lessgtr} = L_{(\alpha_1 + \alpha_2, \max(\beta_1, \beta_2))}^{\lessgtr} \quad (8)$$

$$L_{\gamma_1}^{\lessgtr} \circ_{T_W} L_{\gamma_2}^{\ll} = L_{(\alpha_2 - \alpha_1 + \min(\beta_1, \beta_2) - \beta_1, \max(\beta_1, \beta_2))}^{\ll} \quad (9)$$

$$L_{\gamma_1}^{\ll} \circ_{T_W} L_{\gamma_2}^{\lessgtr} = L_{(\alpha_1 - \alpha_2 + \min(\beta_1, \beta_2) - \beta_2, \max(\beta_1, \beta_2))}^{\lessgtr} \quad (10)$$

where $\gamma_1 = (\alpha_1, \beta_1)$ and $\gamma_2 = (\alpha_2, \beta_2)$.

For $\alpha = 0$, we obtain the following interesting corollary.

Corollary 1: For $\beta \geq 0$ and a, b , and c in \mathbb{R}

$$T_W(L_{(0, \beta)}^{\ll}(a, b), L_{(0, \beta)}^{\ll}(b, c)) \leq L_{(0, \beta)}^{\ll}(a, c) \quad (11)$$

$$T_W(L_{(0, \beta)}^{\lessgtr}(a, b), L_{(0, \beta)}^{\lessgtr}(b, c)) \leq L_{(0, \beta)}^{\lessgtr}(a, c) \quad (12)$$

$$T_W(L_{(0, \beta)}^{\lessgtr}(a, b), L_{(0, \beta)}^{\ll}(b, c)) \leq L_{(0, \beta)}^{\ll}(a, c) \quad (13)$$

$$T_W(L_{(0, \beta)}^{\ll}(a, b), L_{(0, \beta)}^{\lessgtr}(b, c)) \leq L_{(0, \beta)}^{\lessgtr}(a, c) \quad (14)$$

(11) expresses that $L_{(0, \beta)}^{\ll}$ is T_W -transitive while (12) says that $L_{(0, \beta)}^{\lessgtr}$ is T_W -transitive. Furthermore, (13) and (14) express a mixed transitivity between $L_{(0, \beta)}^{\ll}$ and $L_{(0, \beta)}^{\lessgtr}$, generalizing that from $a \leq b$ and $b < c$ it follows that $a < c$, and similarly, that from $a < b$ and $b \leq c$ it follows that $a < c$. Corollary 1 and hence Proposition 1 do not hold for an arbitrary triangular norm T in general.

For $\alpha \geq 0$, $L_{(\alpha, \beta)}^{\lessgtr}$ is reflexive and $L_{(\alpha, \beta)}^{\ll}$ is irreflexive. The following corollary results from the obvious reflexivity and symmetry of $E_{(\alpha, \beta)}$, and Corollary 1.

Corollary 2: $E_{(0, \beta)}$ is a fuzzy T_W -equivalence relation.

From (4) we obtain the T_W - $E_{(0, \beta)}$ -antisymmetry of $L_{(0, \beta)}^{\lessgtr}$. Combined with the reflexivity of $E_{(0, \beta)}$, we establish yet another interesting corollary.

Corollary 3: $L_{(0, \beta)}^{\lessgtr}$ is a fuzzy T_W - $E_{(0, \beta)}$ -ordering.

TABLE II

RELATION BETWEEN THE BOUNDARIES OF THE CRISP INTERVALS $[a^-, a^+]$ AND $[b^-, b^+]$, AND THE FUZZY INTERVALS A AND B .

Crisp	Fuzzy
$a^- < b^-$	$A \circ (L_{(\alpha, \beta)}^{\ll} \triangleright B)$
$a^- \leq b^-$	$(A \circ L_{(\alpha, \beta)}^{\lessgtr}) \triangleright B$
$a^+ < b^+$	$(A \triangleleft L_{(\alpha, \beta)}^{\lessgtr}) \circ B$
$a^+ \leq b^+$	$A \triangleleft (L_{(\alpha, \beta)}^{\lessgtr} \circ B)$
$a^+ < b^-$	$A \triangleleft L_{(\alpha, \beta)}^{\ll} \triangleright B$
$a^+ \leq b^-$	$A \triangleleft L_{(\alpha, \beta)}^{\lessgtr} \triangleright B$
$a^- < b^+$	$A \circ L_{(\alpha, \beta)}^{\lessgtr} \circ B$
$a^- \leq b^+$	$A \circ L_{(\alpha, \beta)}^{\lessgtr} \circ B$

The following proposition is a generalization of the trichotomy law, stating that if b is long before a , a and b cannot be at approximately the same time and a cannot be before b .

Proposition 2: For $0 \leq \beta_1 \leq \beta_2$ and $\alpha_1 \leq \alpha_2$, it holds that

$$T_W(L_{(\alpha_1, \beta_1)}^{\lessgtr}(a, b), L_{(\alpha_2, \beta_2)}^{\ll}(b, a)) = 0 \quad (15)$$

for all a and b in \mathbb{R} .

IV. FUZZY TEMPORAL INTERVAL RELATIONS

A. Ordering of vague boundaries

We define a *fuzzy time period* as a normalised fuzzy set A in \mathbb{R} which is interpreted as the time span of some event. Recall that a fuzzy set A is called normalised if there exists a u in \mathbb{R} such that $A(u) = 1$. Furthermore, a *fuzzy (time) interval* A is a convex and upper semicontinuous normalised fuzzy set in \mathbb{R} , i.e. for each α in $]0, 1]$ the α -level set $\{u | A(u) \geq \alpha\}$ is a closed interval¹. If A and B are fuzzy time intervals, the boundaries of A and B can be gradual. Hence, we cannot refer to these boundaries in the same way we refer to the boundaries of crisp intervals to define temporal relations in the manner of Table I. Nonetheless, we can use the fuzzy orderings between time points defined in the previous section. One possibility to measure for example the extent to which the beginning of a fuzzy time interval A is long before the beginning of a fuzzy time interval B is to look at the highest extent to which there exists a time point in A that occurs long before all time points in B . Similarly, for instance, to express the degree to which the beginning of A is before or at the same time as the ending of B , we can use the highest extent to which there exists a time point in A that occurs before or at the same time as some time point in B . This can be accomplished by using relatedness measures, as shown in Table II. For an arbitrary fuzzy relation R in \mathbb{R} , and fuzzy sets A and B in \mathbb{R} , these

¹All the properties of the fuzzy temporal interval relations in this paper are valid for arbitrary fuzzy time periods, hence from a syntactic point of view neither convexity nor upper semicontinuity are required. However from a semantic point of view, it seems natural to consider only temporal interval relations between fuzzy time intervals since the convexity condition is needed to adequately generalize the notion of an interval, while the upper semicontinuity condition reflects the fact that time intervals are closed intervals.

relatedness measures are defined as [28]:

$$A \circ_T R \circ_T B = \sup_{v \in \mathbb{R}} T(B(v), \sup_{u \in \mathbb{R}} T(A(u), R(u, v))) \quad (16)$$

$$A \triangleleft_I R \triangleright_I B = \inf_{v \in \mathbb{R}} I(B(v), \inf_{u \in \mathbb{R}} I(A(u), R(u, v))) \quad (17)$$

$$(A \triangleleft_I R) \circ_T B = \sup_{v \in \mathbb{R}} T(B(v), \inf_{u \in \mathbb{R}} I(A(u), R(u, v))) \quad (18)$$

$$A \triangleleft_I (R \circ_T B) = \inf_{u \in \mathbb{R}} I(A(u), \sup_{v \in \mathbb{R}} T(B(v), R(u, v))) \quad (19)$$

$$(A \circ_T R) \triangleright_I B = \inf_{v \in \mathbb{R}} I(B(v), \sup_{u \in \mathbb{R}} T(A(u), R(u, v))) \quad (20)$$

$$A \circ_T (R \triangleright_I B) = \sup_{u \in \mathbb{R}} T(A(u), \inf_{v \in \mathbb{R}} I(B(v), R(u, v))) \quad (21)$$

where T is a left-continuous t-norm and I its residual implicator, defined for all x and y in $[0, 1]$ as

$$I(x, y) = \sup\{\lambda \mid T(x, \lambda) \leq y\}$$

These definitions are closely related to the sup- T composition of fuzzy relations, and to the subproduct and superproduct of fuzzy relations [5]. In the remainder of this paper, we will mainly use the Łukasiewicz t-norm T_W and its residual implicator I_W in the definition of the relatedness measures, i.e.

$$I_W(x, y) = \min(1, 1 - x + y)$$

for all x and y in $[0, 1]$. When $T = T_W$ and $I = I_W$, we omit the subscripts of \circ , \triangleleft and \triangleright in (16)–(21). In the remainder of this section, let R and S be fuzzy relations in \mathbb{R} , and A , B , and C fuzzy sets in \mathbb{R} . We recall the following three propositions from [28].

Proposition 3: [28] If A and B are normalised then

$$A \triangleleft (R \circ B) \leq A \circ R \circ B \quad (22)$$

$$(A \circ R) \triangleright B \leq A \circ R \circ B \quad (23)$$

$$A \triangleleft R \triangleright B \leq (A \triangleleft R) \circ B \quad (24)$$

$$A \triangleleft R \triangleright B \leq A \circ (R \triangleright B) \quad (25)$$

$$(A \triangleleft R) \circ B \leq A \triangleleft (R \circ B) \quad (26)$$

$$A \circ (R \triangleright B) \leq (A \circ R) \triangleright B \quad (27)$$

Proposition 4 (Reflexivity): [28] If R is reflexive then

$$A \triangleleft (R \circ A) = 1 \quad (28)$$

$$(A \circ R) \triangleright A = 1 \quad (29)$$

Proposition 5 (Irreflexivity): [28] If R is irreflexive then

$$(A \triangleleft R) \circ A = 0 \quad (30)$$

$$A \circ (R \triangleright A) = 0 \quad (31)$$

Substituting fuzzy time intervals for A and B and either $L_{(\alpha, \beta)}^{\lessdot}$ or $L_{(\alpha, \beta)}^{\ll}$ for R in the propositions above, shows that our approach for modelling relations between the vague boundaries of fuzzy time intervals is sound. For example, from (23) and (27) we derive

$$A \circ (L_{(\alpha, \beta)}^{\ll} \triangleright B) \leq A \circ L_{(\alpha, \beta)}^{\ll} \circ B$$

hence that the degree to which the beginning of A is long before the end of B is at least as high as the degree to which the beginning of A is long before the beginning of B . Furthermore, if $\alpha \geq 0$ to ensure the reflexivity of $L_{(\alpha, \beta)}^{\lessdot}$, from (28) and (29) we derive

$$A \triangleleft (L_{(\alpha, \beta)}^{\lessdot} \circ A) = 1$$

$$(A \circ L_{(\alpha, \beta)}^{\lessdot}) \triangleright A = 1$$

hence that the ending of A is less than or approximately equal to the ending of A to degree 1, and that the beginning of A is less than or approximately equal to the beginning of A to degree 1. In the same way, from (30) and (31) we obtain

$$(A \triangleleft L_{(\alpha, \beta)}^{\ll}) \circ A = 0$$

$$A \circ (L_{(\alpha, \beta)}^{\ll} \triangleright A) = 0$$

in other words that the ending of A is not ‘long before’ the ending of A , and that the beginning of A is not ‘long before’ the beginning of A . Finally, as a result of the following important proposition, the transitivity behaviour of the ordering of the interval boundaries is preserved.

Proposition 6 (Transitivity): For normalised fuzzy sets A , B , and C , the relatedness measures exhibit the transitivity properties displayed in Table III. Let $\mathcal{R}_{(M_1, M_2)}$ be the entry in this table on the row corresponding with the relatedness measure M_1 , and the column corresponding with the relatedness measure M_2 , then it holds that $T_W(M_1, M_2) \leq \mathcal{R}_{(M_1, M_2)}$.

For example, the entry on the sixth line and the third column of Table III should be read as

$$T_W(A \circ (R \triangleright B), (B \triangleleft S) \circ C) \leq A \circ (R \circ S) \circ C$$

Using Proposition 1 we obtain as a special case

$$T_W(A \circ (L_{(0,0)}^{\ll} \triangleright B), (B \triangleleft L_{(0,0)}^{\ll}) \circ C) \leq A \circ L_{(0,0)}^{\ll} \circ C$$

which generalizes the statement that if the beginning of A is before the beginning of B and the ending of B is before the ending of C , then the beginning of A is before the ending of C . This correspondence with the transitivity behaviour of the crisp relations $<$ and \leq also reveals why for some entries in Table III we have no information at all, i.e., the entries that equal 1. For example, from the fact that the beginning of A is before the ending of B , and the fact that the beginning of B is before the ending of C , we can conclude nothing about the relative positioning of A and C . As a consequence, the entry on the first row, first column equals 1.

All results from this section can easily be generalized to an arbitrary universe and an arbitrary left-continuous t-norm. Our commitment to the Łukasiewicz t-norm is mainly motivated by the rich interactions of T_W with $L_{(\alpha, \beta)}^{\ll}$ and $L_{(\alpha, \beta)}^{\lessdot}$, as exemplified by Proposition 1 and Proposition 2.

B. Relations between fuzzy time periods

Using the expressions in Table II it is straightforward to generalize the temporal interval relations from Table I: using the minimum to generalize the conjunctions in Table I, we obtain the generalized definitions in Table IV. Due to the idempotency of the minimum, using the minimum to combine the different constraints on the vague boundaries in this way seems much more natural than, for example, using the Łukasiewicz t-norm. Moreover, it turns out that this choice of the minimum is a prerequisite for some desirable properties of the fuzzy temporal interval relations which will be introduced further on in this section.

TABLE III
TRANSITIVITY TABLE FOR RELATEDNESS MEASURES.

	$B \circ S \circ C$	$B \triangleleft S \triangleright C$	$(B \triangleleft S) \circ C$
$A \circ R \circ B$	1	$A \circ ((R \circ S) \triangleright C)$	$A \circ (R \circ S) \circ C$
$A \triangleleft R \triangleright B$	$(A \triangleleft (R \circ S)) \circ C$	$A \triangleleft (R \circ S) \triangleright C$	$(A \triangleleft (R \circ S)) \circ C$
$(A \triangleleft R) \circ B$	1	$A \triangleleft (R \circ S) \triangleright C$	$(A \triangleleft (R \circ S)) \circ C$
$A \triangleleft (R \circ B)$	1	$A \triangleleft (R \circ S) \triangleright C$	$(A \triangleleft (R \circ S)) \circ C$
$(A \circ R) \triangleright B$	$A \circ (R \circ S) \circ C$	$A \circ ((R \circ S) \triangleright C)$	$A \circ (R \circ S) \circ C$
$A \circ (R \triangleright B)$	$A \circ (R \circ S) \circ C$	$A \circ ((R \circ S) \triangleright C)$	$A \circ (R \circ S) \circ C$

	$B \triangleleft (S \circ C)$	$(B \circ S) \triangleright C$	$B \circ (S \triangleright C)$
$A \circ R \circ B$	$A \circ (R \circ S) \circ C$	1	1
$A \triangleleft R \triangleright B$	$(A \triangleleft (R \circ S)) \circ C$	$A \triangleleft (R \circ S) \triangleright C$	$A \triangleleft (R \circ S) \triangleright C$
$(A \triangleleft R) \circ B$	$(A \triangleleft (R \circ S)) \circ C$	1	1
$A \triangleleft (R \circ B)$	$A \triangleleft ((R \circ S) \circ C)$	1	1
$(A \circ R) \triangleright B$	$A \circ (R \circ S) \circ C$	$(A \circ (R \circ S)) \triangleright C$	$A \circ ((R \circ S) \triangleright C)$
$A \circ (R \triangleright B)$	$A \circ (R \circ S) \circ C$	$A \circ ((R \circ S) \triangleright C)$	$A \circ ((R \circ S) \triangleright C)$

TABLE IV

FUZZY TEMPORAL INTERVAL RELATIONS.


$$\gamma_{123} = (\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3), \gamma_{12} = (\alpha_1, \beta_1, \alpha_2, \beta_2), \gamma_1 = (\alpha_1, \beta_1),$$

$$\gamma_2 = (\alpha_2, \beta_2) \text{ AND } \gamma_3 = (\alpha_3, \beta_3)$$

Name	Notation	Definition
before	$b_{\gamma_1}(A, B)$	$A \triangleleft L_{\gamma_1}^{\leq} \triangleright B$
overlaps	$o_{\gamma_{123}}(A, B)$	$\min(A \circ (L_{\gamma_1}^{\leq} \triangleright B), B \circ L_{\gamma_2}^{\leq} \circ A, (A \triangleleft L_{\gamma_3}^{\leq}) \circ B)$
during	$d_{\gamma_{12}}(A, B)$	$\min(B \circ (L_{\gamma_1}^{\leq} \triangleright A), (A \triangleleft L_{\gamma_2}^{\leq}) \circ B)$
meets	$m_{\gamma_1}(A, B)$	$\min(A \triangleleft L_{\gamma_1}^{\leq} \triangleright B, B \circ L_{\gamma_1}^{\leq} \circ A)$
starts	$s_{\gamma_{12}}(A, B)$	$\min((A \circ L_{\gamma_1}^{\leq}) \triangleright B, (B \circ L_{\gamma_1}^{\leq}) \triangleright A, (A \triangleleft L_{\gamma_2}^{\leq}) \circ B)$
finishes	$f_{\gamma_{12}}(A, B)$	$\min(A \triangleleft (L_{\gamma_1}^{\leq} \circ B), B \triangleleft (L_{\gamma_1}^{\leq} \circ A), B \circ (L_{\gamma_2}^{\leq} \triangleright A))$
equals	$e_{\gamma_{12}}(A, B)$	$\min((A \circ L_{\gamma_1}^{\leq}) \triangleright B, (B \circ L_{\gamma_1}^{\leq}) \triangleright A, A \triangleleft (L_{\gamma_2}^{\leq} \circ B), B \triangleleft (L_{\gamma_2}^{\leq} \circ A))$

Note that the definitions in Table IV coincide with Allen's original definitions if each α_i and β_i equals 0, and A and B are crisp sets. Quantitative information (A happened at least 4 years after B) and semi-quantitative information (A happened long after B) can be expressed using values α_i or β_i different from 0. The (semi-)quantitative information we may have at our disposal about the relative positioning of the beginnings of A and B is independent of the semi-quantitative information we may have at our disposal concerning the endings of A and B ; hence, the fuzzy relation $d_{\gamma_{12}}(A, B)$ involves two different sets of parameters: $\gamma_1 = (\alpha_1, \beta_1)$ and $\gamma_2 = (\alpha_2, \beta_2)$. On the other hand, the two relatedness measures in the definition of $m_{\gamma_1}(A, B)$ together express that the ending of A is approximately equal to the beginning of B , hence the same set of parameters $\gamma_1 = (\alpha_1, \beta_1)$ is used twice. Notice how the notion of approximate equality in the definition of $m_{\gamma_1}(A, B)$ is expressed entirely analogous as in the definition of $E_{(\alpha, \beta)}$ in (4), by making use of $L_{(\alpha_1, \beta_1)}^{\leq}$.

Example 2: For the fuzzy time intervals A and B displayed in Figure 2 it holds that $b_{(0,0)}(A, B) = 1$ and $m_{(0,0)}(A, B) = 0$. In other words, A is considered to be fully before B , as the overlap between A and B is too low for $m(A, B)$ to hold to a degree higher than 0. However, it is clear that A also more or

Fig. 2. Fuzzy time intervals A and B

less meets B . By increasing the value of α we apply a stricter definition of 'long before', and a more tolerant definition of 'approximately at the same time', hence the degree to which A is long before B decreases and the degree to which A more or less meets B increases; we obtain

$$\begin{aligned} b_{(5,0)}(A, B) &= 0.67 & m_{(5,0)}(A, B) &= 0.33 \\ b_{(10,0)}(A, B) &= 0.33 & m_{(10,0)}(A, B) &= 0.67 \\ b_{(15,0)}(A, B) &= 0 & m_{(15,0)}(A, B) &= 1 \end{aligned}$$

When α is sufficiently large, the end of A is not considered to be long before the beginning of B anymore, hence $b_{(15,0)}(A, B) = 0$. A similar observation can be made when increasing the value of β :

$$\begin{aligned} b_{(0,20)}(A, B) &= 0.75 & m_{(0,20)}(A, B) &= 0.25 \\ b_{(0,30)}(A, B) &= 0.5 & m_{(0,30)}(A, B) &= 0.5 \end{aligned}$$

Our generalization preserves several interesting properties of Allen's original algebra, many of which are lost in other approaches. First of all, Allen's temporal interval relations are jointly exhaustive, which means that between any two time intervals at least one of the temporal relations holds. For fuzzy time periods we obtain a generalization, using the Łukasiewicz t -conorm S_W defined for all x and y in $[0, 1]$ as

$$S_W(x, y) = \min(x + y, 1)$$

Proposition 7 (Exhaustivity): Let A and B be fuzzy time

periods. It holds that

$$\begin{aligned} S_W(b_{\gamma_1}(A, B), bi_{\gamma_1}(A, B), o_{\gamma_3}(A, B), oi_{\gamma_3}(A, B), \\ d_{\gamma_2}(A, B), di_{\gamma_2}(A, B), m_{\gamma_1}(A, B), mi_{\gamma_1}(A, B), \\ s_{\gamma_2}(A, B), si_{\gamma_2}(A, B), f_{\gamma_2}(A, B), fi_{\gamma_2}(A, B), \\ e_{\gamma_2}(A, B)) = 1 \end{aligned} \quad (32)$$

For nondegenerate time intervals, i.e. time intervals $[a^-, a^+]$ with $a^- < a^+$, Allen's relations are mutually exclusive. This means that at most one of the temporal relations holds between two given nondegenerate time intervals, and hence precisely one. We call a fuzzy time period A nondegenerate w.r.t. (α, β) iff $A \circ L_{(\alpha, \beta)}^{\ll} \circ A = 1$, i.e., if the beginning of A is long before the end of A . Again, we obtain a generalization of this property:

Proposition 8 (Mutual Exclusiveness): Let A and B be nondegenerate fuzzy time periods w.r.t. $(2\alpha, \beta)$. Moreover, let R and S both be one of the 13 fuzzy temporal relations defined in Table IV, $\alpha_1 = \alpha_2 = \alpha_3 = \alpha \geq 0$ and $\beta_1 = \beta_2 = \beta_3 = \beta \geq 0$. If $R \neq S$ it holds that

$$T_W(R(A, B), S(A, B)) = 0 \quad (33)$$

The condition that A and B should be nondegenerate fuzzy time periods is only needed when R or S is $m_{(\alpha, \beta)}$ or $mi_{(\alpha, \beta)}$. This is not different from the traditional crisp case. For example, using Allen's definitions we have for two crisp intervals $A = [a, b]$ and $B = [b, c]$ that $m(A, B)$ holds. However, if $a = b$, we also have that $s(A, B)$ holds. Likewise, if $b = c$, we have that $f(A, B)$ holds.

Finally, we obtain generalizations of the (a)symmetry and the (ir)reflexivity properties of Allen's relations.

Proposition 9 ((A)symmetry): Let γ_1, γ_{12} and γ_{123} be defined as in Table IV, and let $\alpha_i \geq 0, \beta_i \geq 0$ ($i \in \{1, 2, 3\}$). The relations $b_{\gamma_1}, bi_{\gamma_1}, o_{\gamma_{123}}, oi_{\gamma_{123}}, d_{\gamma_{12}}, di_{\gamma_{12}}, s_{\gamma_{12}}, si_{\gamma_{12}}, f_{\gamma_{12}}$ and $fi_{\gamma_{12}}$ are T_W -asymmetric, i.e., let R be one of the aforementioned fuzzy relations and let A and B be fuzzy time periods. It holds that

$$T_W(R(A, B), R(B, A)) = 0 \quad (34)$$

Furthermore, it holds that

$$e_{\gamma_{12}}(A, B) = e_{\gamma_{12}}(B, A) \quad (35)$$

If A and B are nondegenerate fuzzy time periods w.r.t. (α_1, β_1) , it holds that

$$T_W(m_{\gamma_1}(A, B), m_{\gamma_1}(B, A)) = 0 \quad (36)$$

$$T_W(mi_{\gamma_1}(A, B), mi_{\gamma_1}(B, A)) = 0 \quad (37)$$

Proposition 10 ((Ir)reflexivity): Let γ_1, γ_{12} and γ_{123} be defined as in Table IV, and let $\alpha_i \geq 0, \beta_i \geq 0$ ($i \in \{1, 2, 3\}$). The relations $b_{\gamma_1}, bi_{\gamma_1}, o_{\gamma_{123}}, oi_{\gamma_{123}}, d_{\gamma_{12}}, di_{\gamma_{12}}, s_{\gamma_{12}}, si_{\gamma_{12}}, f_{\gamma_{12}}$ and $fi_{\gamma_{12}}$ are irreflexive, i.e., let R be one of the aforementioned fuzzy relations and let A and B be fuzzy time periods. It holds that

$$R(A, A) = 0 \quad (38)$$

Furthermore, it holds that

$$e_{\gamma_{12}}(A, A) = 1 \quad (39)$$

If A is a nondegenerate fuzzy time period w.r.t. (α_1, β_1) , it holds that

$$m_{\gamma_1}(A, A) = mi_{\gamma_1}(A, A) = 0 \quad (40)$$

In Proposition 7–10, fuzzy relations of the form $L_{(\alpha, \beta)}^{\ll}$ and $L_{(\alpha, \beta)}^{\lessdot}$ are used to express the concepts 'long before' and 'more or less before'. In principle, more general classes of fuzzy relations could be used to this end, i.e., fuzzy relations that cannot be written as either $L_{(\alpha, \beta)}^{\ll}$ or $L_{(\alpha, \beta)}^{\lessdot}$. However, as can easily be seen from their proof in Appendix III, these propositions remain valid for more general classes of fuzzy relations, provided some weak assumptions are satisfied. For example, let R and S be arbitrary fuzzy relations in \mathbb{R} which are used to express the concepts 'long before' and 'more or less before' respectively. Then, Proposition 7 remains valid if $R(x, y) = 1 - S(y, x)$ for all x and y in \mathbb{R} . For Proposition 8–10 to hold, we also have to assume, among others, that $R, R \circ S, R \circ S \circ S$, etc. are irreflexive.

However, using fuzzy relations of the form $L_{(\alpha, \beta)}^{\ll}$ and $L_{(\alpha, \beta)}^{\lessdot}$ to express fuzzy orderings of time points, has a number of important advantages. As shown in Section III-B, these fuzzy relations satisfy many desirable properties, and their sup- T_W composition can be conveniently characterized (Proposition 1), which is important for reasoning with fuzzy temporal relations. Moreover, in [30], we have shown that this choice allows to evaluate the fuzzy temporal interval relations in an efficient way for piecewise linear fuzzy intervals, an important prerequisite for most real-world applications.

V. FUZZY TEMPORAL REASONING

When $A = [a^-, a^+]$, $B = [b^-, b^+]$ and $C = [c^-, c^+]$ are crisp intervals, using Allen's original definitions we can deduce, for example, from $d(A, B)$ and $m(B, C)$ that $b(A, C)$ holds. Indeed by $d(A, B)$ we have $a^+ < b^+$ and by $m(B, C)$ we have $b^+ = c^-$; from $a^+ < b^+$ and $b^+ = c^-$ we conclude $a^+ < c^-$, or in other words $b(A, C)$. When A, B and C are fuzzy time intervals, we would like to make similar deductions, even when the interval relations are imprecise (i.e., $\alpha_i > 0$ or $\beta_i > 0$). To this end, we use the Łukasiewicz t-norm T_W to generalize such deductions. For example, let A, B and C be fuzzy time intervals, $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ and $\beta_1, \beta_2, \beta_3 \geq 0$. Furthermore let $\gamma_{12} = (\alpha_1, \beta_1, \alpha_2, \beta_2)$, $\gamma_1 = (\alpha_1, \beta_1)$, $\gamma_2 = (\alpha_2, \beta_2)$ and $\gamma_3 = (\alpha_3, \beta_3)$ as before. We obtain

$$\begin{aligned} T_W(d_{\gamma_{12}}(A, B), m_{\gamma_3}(B, C)) \\ = T_W(\min(B \circ (L_{\gamma_1}^{\ll} \triangleright A), (A \triangleleft L_{\gamma_2}^{\ll}) \circ B), \\ \min(B \triangleleft L_{\gamma_3}^{\lessdot} \triangleright C, C \circ L_{\gamma_3}^{\lessdot} \circ B)) \\ \leq \min(T_W(B \circ (L_{\gamma_1}^{\ll} \triangleright A), B \triangleleft L_{\gamma_3}^{\lessdot} \triangleright C), \\ T_W(B \circ (L_{\gamma_1}^{\ll} \triangleright A), C \circ L_{\gamma_3}^{\lessdot} \circ B), \\ T_W((A \triangleleft L_{\gamma_2}^{\ll}) \circ B, B \triangleleft L_{\gamma_3}^{\lessdot} \triangleright C), \\ T_W((A \triangleleft L_{\gamma_2}^{\ll}) \circ B, C \circ L_{\gamma_3}^{\lessdot} \circ B)) \\ \leq \min(T_W(B \circ (L_{\gamma_1}^{\ll} \triangleright A), C \circ L_{\gamma_3}^{\lessdot} \circ B), \\ T_W((A \triangleleft L_{\gamma_2}^{\ll}) \circ B, B \triangleleft L_{\gamma_3}^{\lessdot} \triangleright C)) \end{aligned}$$

Using Table III, i.e., the transitivity table for relatedness measures and (10), we obtain

$$\begin{aligned} &\leq \min(1, A \triangleleft (L_{\gamma_2}^{\ll} \circ L_{\gamma_3}^{\ll}) \triangleright C) \\ &= A \triangleleft (L_{\gamma_2}^{\ll} \circ L_{\gamma_3}^{\ll}) \triangleright C \\ &= A \triangleleft L_{\gamma}^{\ll} \triangleright C \\ &= b_{\gamma}(A, C) \end{aligned}$$

where $\gamma = (\alpha_2 - \alpha_3 + \min(\beta_2, \beta_3) - \beta_3, \max(\beta_2, \beta_3))$. In particular when $\alpha_1 = \alpha_2 = \alpha_3 = 0$ and $\beta_1 = \beta_2 = \beta_3 = \beta$ we have $T_W(d_{(0,\beta,0,\beta)}(A, B), m_{(0,\beta)}(B, C)) \leq b_{(0,\beta)}(A, C)$, stating how the degree to which A is during B and the degree to which B more or less meets C , can be used to compute a lower bound for the degree to which A is long before C . This is a generalization of the statement that if A occurs during B and B meets C , then A occurs before C .

As another example, in the crisp case from $m(A, B)$ and $m(B, C)$ one can conclude that $b(A, C)$ holds, under the assumption that B is a nondegenerate interval [3]. In our generalized approach we obtain for fuzzy time intervals A , B and C

$$\begin{aligned} &T_W(m_{(0,\beta)}(A, B), m_{(0,\beta)}(B, C)) \\ &= T_W(\min(A \triangleleft L_{(0,\beta)}^{\ll} \triangleright B, B \circ L_{(0,\beta)}^{\ll} \circ A), \\ &\quad \min(B \triangleleft L_{(0,\beta)}^{\ll} \triangleright C, C \circ L_{(0,\beta)}^{\ll} \circ B)) \\ &\leq T_W(A \triangleleft L_{(0,\beta)}^{\ll} \triangleright B, B \triangleleft L_{(0,\beta)}^{\ll} \triangleright C) \end{aligned}$$

Assuming that B is nondegenerate w.r.t. $(0, \beta)$, i.e. that $B \circ L_{(0,\beta)}^{\ll} \circ B = 1$, and using Table III and (9), we obtain

$$\begin{aligned} A \triangleleft L_{(0,\beta)}^{\ll} \triangleright B &= T_W(A \triangleleft L_{(0,\beta)}^{\ll} \triangleright B, B \circ L_{(0,\beta)}^{\ll} \circ B) \\ &\leq (A \triangleleft (L_{(0,\beta)}^{\ll} \circ L_{(0,\beta)}^{\ll})) \circ B \\ &= (A \triangleleft L_{(0,\beta)}^{\ll}) \circ B \end{aligned}$$

and thus, using Table III and (10),

$$\begin{aligned} &T_W(A \triangleleft L_{(0,\beta)}^{\ll} \triangleright B, B \triangleleft L_{(0,\beta)}^{\ll} \triangleright C) \\ &\leq T_W((A \triangleleft L_{(0,\beta)}^{\ll}) \circ B, B \triangleleft L_{(0,\beta)}^{\ll} \triangleright C) \\ &\leq A \triangleleft (L_{(0,\beta)}^{\ll} \circ L_{(0,\beta)}^{\ll}) \triangleright C \\ &= A \triangleleft L_{(0,\beta)}^{\ll} \triangleright C \\ &= b_{(0,\beta)}(A, C) \end{aligned}$$

This deduction process can easily be automated, which is what we have done to obtain Table V. In the crisp case, the temporal relation that results from composing two temporal relations is not always fully determined. For example, for crisp intervals A , B and C such that $m(A, B)$ and $d(B, C)$, we have that $o(A, C)$, $d(A, C)$ or $s(A, C)$ may hold since we can deduce only that $a^+ > c^-$ and $a^+ < c^+$. Freksa [13] defined a set of coarser temporal relations, which he calls conceptual neighborhoods, and provided a transitivity table that is deductively closed for Allen's original relations as well as the conceptual neighborhoods. For example $o(A, C)$ or $d(A, C)$ or $s(A, C)$ is equivalent to $bc(A, C)$. The definitions of the relevant conceptual neighborhoods are shown in Table VI. Generalizing these definitions to cope with fuzzy time intervals and

imprecise temporal relations is straightforward, using again the relatedness measures from Table II. To obtain Table V, we have assumed that A , B and C are nondegenerate fuzzy time intervals w.r.t. $(0, 0)$. One can verify that when A , B and C are crisp intervals, Table V corresponds to Freksa's transitivity table. As a consequence, by restricting Table V to the first 13 rows and the first 13 columns, we obtain a transitivity table which is a sound generalization of Allen's transitivity table. Note that while Table III serves to derive knowledge about relationships between the gradual boundaries of fuzzy intervals, Table V is used to reason about the relationships between fuzzy intervals themselves.

Table V cannot be used for reasoning with (semi-) quantitative temporal information, i.e., when some $\alpha_i \neq 0$ or $\beta_i \neq 0$. It is not feasible to construct a more general transitivity table, which would permit this and which is still deductively closed. Instead, when $\alpha \neq 0$ or $\beta \neq 0$, the transitivity table for relatedness measures can be used to make deductions. For example, it holds that

$$\begin{aligned} &T_W(o_{\gamma_{123}}(A, B), d_{\gamma_{45}}(B, C)) \\ &\leq \min(C \circ L_{\gamma'}^{\ll} \circ A, A \circ L_{\gamma''}^{\ll} \circ C, \\ &\quad (A \triangleleft L_{\gamma'''}^{\ll}) \circ C) \end{aligned}$$

where γ_{123} is defined as in Table IV and

$$\begin{aligned} \gamma_{45} &= (\alpha_4, \beta_4, \alpha_5, \beta_5) \\ \gamma &= (\alpha_2 + \alpha_4 + \min(\beta_2, \beta_4), \max(\beta_2, \beta_4)) \\ \gamma' &= (\alpha_1 + \alpha_5 + \min(\beta_1, \beta_5), \max(\beta_1, \beta_5)) \\ \gamma'' &= (\alpha_3 + \alpha_5 + \min(\beta_3, \beta_5), \max(\beta_3, \beta_5)) \end{aligned}$$

This result can only be written as $bc_{\gamma''''}$ for a given γ'''' if

$$(A \triangleleft L_{\gamma''''}^{\ll}) \circ C \leq A \circ L_{\gamma'}^{\ll} \circ C$$

which does not hold for arbitrary α_i and β_i .

Note that a lot of interesting theoretical results related to temporal reasoning with crisp intervals have already been established (e.g., [18], [31], [32]). Among others, these results pertain to constructing maximal subalgebras of Allen's original algebra for which reasoning with a composition table is sufficient to detect inconsistencies (i.e., satisfiability), and to find all consequences of an initial set of assumptions (i.e., completeness). Generalizing these results, however, is clearly outside the scope of this paper.

VI. APPLICATION TO QUESTION ANSWERING

The list question "Who was president of the USA during the Cold War?" is problematic for a traditional QA-system, since there is no generally accepted starting date and ending date for the Cold War. Assume however that, while searching the web, our QA-system has discovered the following relevant statements:

- Truman took office when President Roosevelt died just before the end of World War II, ... ²

²<http://www.juntosociety.com/uspresidents/hstruman.html> (accessed June 30, 2005)

TABLE V

TRANSITIVITY TABLE FOR FUZZY TEMPORAL INTERVAL RELATIONS ($\gamma^1 = (0, 0), \gamma^2 = (0, 0, 0, 0), \gamma^3 = (0, 0, 0, 0, 0, 0)$). LET $\mathcal{R}_{(R_1, R_2)}$ BE THE ENTRY IN THIS TABLE ON THE ROW CORRESPONDING WITH THE FUZZY TEMPORAL RELATION R_1 AND THE COLUMN CORRESPONDING WITH THE FUZZY TEMPORAL RELATION R_2 . FOR NONDEGENERATE (W.R.T. $(0, 0)$) FUZZY TIME PERIODS A, B AND C , IT HOLDS THAT

$$T_W(R_1(A, B), R_2(B, C)) \leq \mathcal{R}_{(R_1, R_2)}(A, C).$$

	b_{γ^1}	m_{γ^1}	o_{γ^3}	fi_{γ^2}	di_{γ^2}	si_{γ^2}	e_{γ^2}	s_{γ^2}	d_{γ^2}	f_{γ^2}	oi_{γ^3}	mi_{γ^1}	bi_{γ^1}	ob_{γ^2}
b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	b_{γ^1}
m_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	b_{γ^1}	m_{γ^1}	m_{γ^1}	m_{γ^1}	bc_{γ^2}	bc_{γ^2}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	b_{γ^1}
o_{γ^3}	b_{γ^1}	b_{γ^1}	ob_{γ^2}	ob_{γ^2}	ol_{γ^1}	oc_{γ^2}	o_{γ^3}	o_{γ^3}	bc_{γ^2}	bc_{γ^2}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	ob_{γ^2}
fi_{γ^2}	b_{γ^1}	m_{γ^1}	o_{γ^3}	fi_{γ^2}	di_{γ^2}	di_{γ^2}	fi_{γ^2}	o_{γ^3}	bc_{γ^2}	tt_{γ^1}	sc_{γ^2}	sc_{γ^2}	sv_{γ^1}	ob_{γ^2}
di_{γ^2}	ol_{γ^1}	oc_{γ^2}	oc_{γ^2}	di_{γ^2}	di_{γ^2}	di_{γ^2}	di_{γ^2}	ct_{γ^2}	ct_{γ^2}	sc_{γ^2}	sc_{γ^2}	sc_{γ^2}	sv_{γ^1}	ol_{γ^1}
si_{γ^2}	ol_{γ^1}	oc_{γ^2}	oc_{γ^2}	di_{γ^2}	di_{γ^2}	si_{γ^2}	si_{γ^2}	hh_{γ^1}	yc_{γ^2}	oi_{γ^3}	oi_{γ^3}	mi_{γ^1}	bi_{γ^1}	ol_{γ^1}
e_{γ^2}	b_{γ^1}	m_{γ^1}	o_{γ^3}	fi_{γ^2}	di_{γ^2}	si_{γ^2}	e_{γ^2}	s_{γ^2}	d_{γ^2}	f_{γ^2}	oi_{γ^3}	mi_{γ^1}	bi_{γ^1}	ob_{γ^2}
s_{γ^2}	b_{γ^1}	b_{γ^1}	ob_{γ^2}	ob_{γ^2}	ol_{γ^1}	hh_{γ^1}	s_{γ^2}	s_{γ^2}	d_{γ^2}	d_{γ^2}	yc_{γ^2}	mi_{γ^1}	bi_{γ^1}	ob_{γ^2}
d_{γ^2}	b_{γ^1}	m_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	yo_{γ^1}	d_{γ^2}	d_{γ^2}	d_{γ^2}	d_{γ^2}	yo_{γ^1}	yo_{γ^1}	bi_{γ^1}	sb_{γ^1}
f_{γ^2}	b_{γ^1}	m_{γ^1}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	ys_{γ^2}	f_{γ^2}	d_{γ^2}	d_{γ^2}	f_{γ^2}	ys_{γ^2}	bi_{γ^1}	bi_{γ^1}	sb_{γ^1}
oi_{γ^3}	ol_{γ^1}	oc_{γ^2}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	ys_{γ^2}	oi_{γ^3}	yc_{γ^2}	yc_{γ^2}	oi_{γ^3}	ys_{γ^2}	bi_{γ^1}	bi_{γ^1}	bd_{γ^1}
mi_{γ^1}	ol_{γ^1}	hh_{γ^1}	yc_{γ^2}	mi_{γ^1}	bi_{γ^1}	bi_{γ^1}	mi_{γ^1}	yc_{γ^2}	yc_{γ^2}	mi_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	bd_{γ^1}
bi_{γ^1}	1	yo_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	yo_{γ^1}	yo_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	1
ob_{γ^2}	b_{γ^1}	b_{γ^1}	ob_{γ^2}	ob_{γ^2}	ol_{γ^1}	ol_{γ^1}	ob_{γ^2}	ob_{γ^2}	sb_{γ^1}	sb_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	ob_{γ^2}
oc_{γ^2}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	oc_{γ^2}	oc_{γ^2}	oc_{γ^2}	ct_{γ^2}	ct_{γ^2}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	ol_{γ^1}
hh_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	hh_{γ^1}	hh_{γ^1}	hh_{γ^1}	yc_{γ^2}	yc_{γ^2}	yc_{γ^2}	mi_{γ^1}	bi_{γ^1}	ol_{γ^1}
yc_{γ^2}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	yo_{γ^1}	yc_{γ^2}	yc_{γ^2}	yc_{γ^2}	yc_{γ^2}	yo_{γ^1}	yo_{γ^1}	bi_{γ^1}	bd_{γ^1}
bc_{γ^2}	b_{γ^1}	b_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	db_{γ^1}	bc_{γ^2}	bc_{γ^2}	bc_{γ^2}	bc_{γ^2}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	sb_{γ^1}
tt_{γ^1}	b_{γ^1}	m_{γ^1}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	sv_{γ^1}	tt_{γ^1}	bc_{γ^2}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	sb_{γ^1}
sc_{γ^2}	ol_{γ^1}	oc_{γ^2}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	sv_{γ^1}	sc_{γ^2}	ct_{γ^2}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	bd_{γ^1}
ys_{γ^2}	1	db_{γ^1}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	ys_{γ^2}	ys_{γ^2}	yo_{γ^1}	yo_{γ^1}	ys_{γ^2}	ys_{γ^2}	sv_{γ^1}	bi_{γ^1}	1
ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	ol_{γ^1}
yo_{γ^1}	1	1	1	1	1	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	bi_{γ^1}	bi_{γ^1}	1
sb_{γ^1}	b_{γ^1}	b_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	1	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	1	1	sb_{γ^1}
sv_{γ^1}	1	db_{γ^1}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	db_{γ^1}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	sv_{γ^1}	1
ct_{γ^2}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	db_{γ^1}	ct_{γ^2}	ct_{γ^2}	ct_{γ^2}	ct_{γ^2}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	bd_{γ^1}
bd_{γ^1}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	1	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	1	1	bd_{γ^1}
db_{γ^1}	1	1	1	1	1	db_{γ^1}	db_{γ^1}	db_{γ^1}	db_{γ^1}	db_{γ^1}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	1

	oc_{γ^2}	hh_{γ^1}	yc_{γ^2}	bc_{γ^2}	tt_{γ^1}	sc_{γ^2}	ys_{γ^2}	ol_{γ^1}	yo_{γ^1}	sb_{γ^1}	sv_{γ^1}	ct_{γ^2}	bd_{γ^1}	db_{γ^1}
b_{γ^1}	b_{γ^1}	b_{γ^1}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	b_{γ^1}	1	sb_{γ^1}	1	sb_{γ^1}	sb_{γ^1}	1
m_{γ^1}	b_{γ^1}	m_{γ^1}	bc_{γ^2}	sb_{γ^1}	sb_{γ^1}	sb_{γ^1}	db_{γ^1}	b_{γ^1}	db_{γ^1}	sb_{γ^1}	1	sb_{γ^1}	sb_{γ^1}	1
o_{γ^3}	ol_{γ^1}	oc_{γ^2}	ct_{γ^2}	sb_{γ^1}	sb_{γ^1}	bd_{γ^1}	db_{γ^1}	ol_{γ^1}	db_{γ^1}	sb_{γ^1}	1	bd_{γ^1}	bd_{γ^1}	1
fi_{γ^2}	oc_{γ^2}	oc_{γ^2}	ct_{γ^2}	bc_{γ^2}	tt_{γ^1}	sc_{γ^2}	sv_{γ^1}	ol_{γ^1}	db_{γ^1}	sb_{γ^1}	sv_{γ^1}	ct_{γ^2}	bd_{γ^1}	db_{γ^1}
di_{γ^2}	oc_{γ^2}	oc_{γ^2}	ct_{γ^2}	ct_{γ^2}	sc_{γ^2}	sc_{γ^2}	sv_{γ^1}	ol_{γ^1}	db_{γ^1}	bd_{γ^1}	sv_{γ^1}	ct_{γ^2}	bd_{γ^1}	db_{γ^1}
si_{γ^2}	oc_{γ^2}	hh_{γ^1}	yc_{γ^2}	ct_{γ^2}	sc_{γ^2}	sc_{γ^2}	ys_{γ^2}	ol_{γ^1}	yo_{γ^1}	bd_{γ^1}	sv_{γ^1}	ct_{γ^2}	bd_{γ^1}	db_{γ^1}
e_{γ^2}	oc_{γ^2}	hh_{γ^1}	yc_{γ^2}	bc_{γ^2}	tt_{γ^1}	sc_{γ^2}	ys_{γ^2}	ol_{γ^1}	yo_{γ^1}	sb_{γ^1}	sv_{γ^1}	ct_{γ^2}	bd_{γ^1}	db_{γ^1}
s_{γ^2}	ol_{γ^1}	hh_{γ^1}	yc_{γ^2}	sb_{γ^1}	sb_{γ^1}	bd_{γ^1}	yo_{γ^1}	ol_{γ^1}	yo_{γ^1}	sb_{γ^1}	1	bd_{γ^1}	bd_{γ^1}	1
d_{γ^2}	1	yo_{γ^1}	yo_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	yo_{γ^1}	1	yo_{γ^1}	sb_{γ^1}	1	1	1	1
f_{γ^2}	db_{γ^1}	yo_{γ^1}	yo_{γ^1}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	ys_{γ^2}	1	yo_{γ^1}	sb_{γ^1}	sv_{γ^1}	db_{γ^1}	1	db_{γ^1}
oi_{γ^3}	db_{γ^1}	yo_{γ^1}	yo_{γ^1}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	ys_{γ^2}	1	yo_{γ^1}	bd_{γ^1}	sv_{γ^1}	db_{γ^1}	1	db_{γ^1}
mi_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	yc_{γ^2}	mi_{γ^1}	bi_{γ^1}	bi_{γ^1}	1	yo_{γ^1}	bd_{γ^1}	bi_{γ^1}	yo_{γ^1}	1	yo_{γ^1}
bi_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	yo_{γ^1}	bi_{γ^1}	bi_{γ^1}	bi_{γ^1}	1	yo_{γ^1}	1	bi_{γ^1}	yo_{γ^1}	1	yo_{γ^1}
ob_{γ^2}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	sb_{γ^1}	sb_{γ^1}	bd_{γ^1}	1	ol_{γ^1}	1	sb_{γ^1}	1	bd_{γ^1}	bd_{γ^1}	1
oc_{γ^2}	ol_{γ^1}	oc_{γ^2}	ct_{γ^2}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	db_{γ^1}	ol_{γ^1}	db_{γ^1}	bd_{γ^1}	1	bd_{γ^1}	bd_{γ^1}	1
hh_{γ^1}	ol_{γ^1}	hh_{γ^1}	yc_{γ^2}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	yo_{γ^1}	ol_{γ^1}	yo_{γ^1}	bd_{γ^1}	1	bd_{γ^1}	bd_{γ^1}	1
yc_{γ^2}	1	yo_{γ^1}	yo_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	yo_{γ^1}	1	yo_{γ^1}	bd_{γ^1}	1	1	1	1
bc_{γ^2}	1	db_{γ^1}	db_{γ^1}	sb_{γ^1}	sb_{γ^1}	1	db_{γ^1}	1	db_{γ^1}	sb_{γ^1}	1	1	1	1
tt_{γ^1}	db_{γ^1}	db_{γ^1}	db_{γ^1}	bc_{γ^2}	tt_{γ^1}	sv_{γ^1}	sv_{γ^1}	1	db_{γ^1}	sb_{γ^1}	sv_{γ^1}	db_{γ^1}	1	db_{γ^1}
sc_{γ^2}	db_{γ^1}	db_{γ^1}	db_{γ^1}	ct_{γ^2}	sc_{γ^2}	sv_{γ^1}	sv_{γ^1}	1	db_{γ^1}	bd_{γ^1}	sv_{γ^1}	db_{γ^1}	1	db_{γ^1}
ys_{γ^2}	db_{γ^1}	yo_{γ^1}	yo_{γ^1}	db_{γ^1}	sv_{γ^1}	sv_{γ^1}	ys_{γ^2}	1	yo_{γ^1}	1	sv_{γ^1}	db_{γ^1}	1	db_{γ^1}
ol_{γ^1}	ol_{γ^1}	ol_{γ^1}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	bd_{γ^1}	1	ol_{γ^1}	1	bd_{γ^1}	1	bd_{γ^1}		

TABLE VI
FREKSA'S CONCEPTUAL NEIGHBORHOODS [13].

Name	Definition
older	$ol(A, B) \equiv a^- < b^-$
head to head with	$hh(A, B) \equiv a^- = b^-$
younger	$yo(A, B) \equiv b^- < a^-$
survived by	$sb(A, B) \equiv a^+ < b^+$
tail to tail with	$tt(A, B) \equiv a^+ = b^+$
survives	$sv(A, B) \equiv b^+ < a^+$
born before death of	$bd(A, B) \equiv a^- < b^+$
contemporary of	$ct(A, B) \equiv a^- < b^+ \text{ and } b^- < a^+$
died after birth of	$db(A, B) \equiv b^- < a^+$
older & survived by	$ob(A, B) \equiv a^- < b^- \text{ and } a^+ < b^+$
older contemporary of	$oc(A, B) \equiv a^- < b^- \text{ and } b^- < a^+$
surviving contemporary of	$sc(A, B) \equiv a^- < b^+ \text{ and } b^+ < a^+$
survived by contemporary of	$bc(A, B) \equiv b^- < a^+ \text{ and } a^+ < b^+$
younger contemporary of	$yc(A, B) \equiv b^- < a^- \text{ and } a^- < b^+$
younger & survives	$ys(A, B) \equiv b^- < a^- \text{ and } b^+ < a^+$

- The Cold War began after World War Two. ³
- The Cold War began at the close of the WW II and ended with the dissolution of the Soviet Union. ⁴
- President (...) George H.W. Bush declared a U.S.-Soviet strategic partnership at the summit of July 1991, decisively marking the end of the Cold War. ⁵

For ease of notation, we use the following abbreviations:

FDR	denotes the life span of Franklin Roosevelt
TRP	denotes the time span of the presidency of Harry Truman
$WW2$	denotes the time span of the second world war
CW	denotes the time span of the cold war
BSP	denotes the time span of the presidency of George H.W. Bush

Dates are treated as real numbers that express the number of years since January 1, 1900. As temporal information extraction is not the focus of this paper, we assume that we have the following interpretation of the above statements at our disposal:

$$\begin{aligned}
m_{(0,0)}(FDR, TRP) &= 1 \\
tt_{(0.3,0.5)}(FDR, WW2) &= 1 \\
sb_{(0,0)}(FDR, WW2) &= 1 \\
b_{(0,0)}(WW2, CW) &= 1 \\
m_{(0.3,0.5)}(WW2, CW) &= 1 \\
o_{(0,0)}(CW, BSP) &= 1
\end{aligned}$$

Note that $(\alpha, \beta) = (0.3, 0.5)$ means that two time points within a period of 3.6 months are considered approximately equal to degree 1, and two time points within a period of 9.6 months are approximately equal to a degree that is higher than

³http://www.globalsecurity.org/military/ops/cold_war.htm (accessed June 30, 2005)

⁴<http://www.videofact.com/coldwar.htm> (accessed June 30, 2005)

⁵[http://www.absoluteastronomy.com/encyclopedia/H/Hi/History_of_the_Soviet_Union_\(1985-1991\).htm](http://www.absoluteastronomy.com/encyclopedia/H/Hi/History_of_the_Soviet_Union_(1985-1991).htm) (accessed June 30, 2005)

0. Using Table III and (8), we obtain

$$\begin{aligned}
&T_W(tt_{(0.3,0.5)}(FDR, WW2), m_{(0.3,0.5)}(WW2, CW)) \\
&= T_W(\min(FDR \triangleleft (L_{(0.3,0.5)}^{\lessdot} \circ WW2), \\
&\quad WW2 \triangleleft (L_{(0.3,0.5)}^{\lessdot} \circ FDR)), \\
&\quad \min(WW2 \triangleleft L_{(0.3,0.5)}^{\lessdot} \triangleright CW, \\
&\quad CW \circ L_{(0.3,0.5)}^{\lessdot} \circ WW2)) \\
&\leq T_W(WW2 \triangleleft (L_{(0.3,0.5)}^{\lessdot} \circ FDR), \\
&\quad CW \circ L_{(0.3,0.5)}^{\lessdot} \circ WW2) \\
&\leq CW \circ (L_{(0.3,0.5)}^{\lessdot} \circ L_{(0.3,0.5)}^{\lessdot}) \circ FDR \\
&= CW \circ L_{(0.6,0.5)}^{\lessdot} \circ FDR
\end{aligned}$$

Furthermore, using Table III and (9) we have,

$$\begin{aligned}
&T_W(CW \circ L_{(0.6,0.5)}^{\lessdot} \circ FDR, m_{(0,0)}(FDR, TRP)) \\
&\leq T_W(CW \circ L_{(0.6,0.5)}^{\lessdot} \circ FDR, \\
&\quad \min(FDR \triangleleft L_{(0,0)}^{\lessdot} \triangleright TRP, \\
&\quad TRP \circ L_{(0,0)}^{\lessdot} \circ FDR)) \\
&\leq T_W(CW \circ L_{(0.6,0.5)}^{\lessdot} \circ FDR, \\
&\quad FDR \triangleleft L_{(0,0)}^{\lessdot} \triangleright TRP) \\
&\leq CW \circ ((L_{(0.6,0.5)}^{\lessdot} \circ L_{(0,0)}^{\lessdot}) \triangleright TRP) \\
&= CW \circ (L_{(0.6,0.5)}^{\lessdot} \triangleright TRP)
\end{aligned}$$

and since we can assume that TRP is nondegenerate w.r.t. $(0.6, 0.5)$ (i.e., Truman was president for more than 13.2 months), we obtain

$$\begin{aligned}
&= T_W(CW \circ (L_{(0.6,0.5)}^{\lessdot} \triangleright TRP), TRP \circ L_{(0.6,0.5)}^{\lessdot} \circ TRP) \\
&\leq CW \circ (L_{(0.6,0.5)}^{\lessdot} \circ L_{(0.6,0.5)}^{\lessdot}) \circ TRP \\
&= CW \circ L_{(0,0.5)}^{\lessdot} \circ TRP \\
&= bd_{(0,0.5)}(CW, TRP)
\end{aligned}$$

From Table V we immediately have

$$\begin{aligned}
&T_W(mi_{(0,0)}(TRP, FDR), sb_{(0,0)}(FDR, WW2)) \\
&\leq bd_{(0,0)}(TRP, WW2) \\
&T_W(bd_{(0,0)}(TRP, WW2), b_{(0,0)}(WW2, CW)) \\
&\leq ol_{(0,0)}(TRP, CW)
\end{aligned}$$

Hence we have that when Harry Truman became president, the Cold War had not yet started, i.e., $ol_{(0,0)}(TRP, CW)$, while it had already started when his presidency ended, i.e., $bd_{(0,0.5)}(CW, TRP)$. In other words, Truman was the first to be president of the USA during the Cold War. Moreover $o_{(0,0)}(CW, BSP)$ means that George H.W. Bush was the last to be president of the USA during the cold war. To answer the question under consideration we only have to determine who was president of the USA between Harry Truman and George H.W. Bush, which is a fairly simple task.

Another solution would be to construct a fuzzy time interval CW which represents the time span of the Cold War. Such a fuzzy time interval is illustrated in Figure 3. For each




Fig. 3. Modelling the time span of the Cold War

president X of the USA, $ct_{(0,0)}(CW, X)$ expresses the degree to which X has been president during the Cold War. A clear advantage of this method is that we can make a difference between presidents that were definitely president during the Cold War (e.g., Richard Nixon, 1969–1974) and presidents whose presidency was more or less during the Cold War (e.g., Harry Truman, 1945–1953). A disadvantage of this method is that the automatic construction of the fuzzy time interval CW is more time-consuming. Moreover, in some situations we may lack sufficient information to construct the fuzzy time interval corresponding to a given imprecise event.

In practice, both approaches may be combined, yielding a system that simply evaluates which fuzzy temporal relations hold if a suitable fuzzy time span can be constructed, and applies fuzzy temporal reasoning otherwise. We refer to [2], [26] for more details on the architecture and implementation of such a system. Currently, only qualitative temporal relations are considered in this system. This is due to the fact that interpreting a natural language statement such as “ X began just before the end of Y ”, i.e., providing suitable values for the parameters α and β , is far from trivial.

VII. CONCLUSIONS

We have suggested a general approach to represent and compute precise and imprecise temporal relations between crisp as well as fuzzy time intervals. To this end, we have used fuzzy orderings of time points, which are lifted into interval relations through the use of relatedness measures. We have shown that both the fuzzy orderings of time points and the relatedness measures satisfy many desirable properties regarding (ir)reflexivity, (a)symmetry and transitivity. When considering only precise relations between crisp time intervals, our approach coincides with Allen’s temporal interval algebra. However, even in the most general model, i.e. for imprecise relations between fuzzy intervals, unlike in previous approaches, generalizations of all the important properties of Allen’s interval relations are valid, in particular those related to exhaustivity, mutual exclusiveness, (ir)reflexivity and (a)symmetry. Moreover, a sound generalization of Freksa’s transitivity table was given which can be used for fuzzy temporal reasoning with qualitative temporal information. In general, fuzzy temporal reasoning can easily be automated by using the transitivity table for relatedness measures which we have introduced in this paper. Finally, we have provided an example which illustrates how fuzzy temporal reasoning

could be useful for (temporal) question answering systems, as these systems have to deal with both imprecise events (e.g., the Cold War) and imprecise temporal relations expressed in natural language (e.g., Roosevelt died just before the end of the Second World War).

APPENDIX I

PROOF OF PROPERTIES CONCERNING THE FUZZY ORDERINGS $L_{(\alpha,\beta)}^{\ll}$ AND $L_{(\alpha,\beta)}^{\gg}$

Throughout the appendices, let T denote a left-continuous t-norm (i.e., a t-norm with left-continuous partial mappings); then T and its residual implicator I satisfy the residuation principle, i.e., for all a, b and c in $[0, 1]$ it holds that

$$T(a, b) \leq c \Leftrightarrow a \leq I(b, c) \quad (41)$$

Moreover, it can be shown that for all a, b, c and d in $[0, 1]$ it holds that (see e.g., [19])

$$T(I(a, b), c) \leq I(a, T(b, c)) \quad (42)$$

$$T(a, I(a, b)) \leq b \quad (43)$$

$$I(T(a, b), c) = I(a, I(b, c)) \quad (44)$$

$$I(a, I_T(b, c)) = I(b, I(a, c)) \quad (45)$$

$$T(I(a, b), I(c, d)) \leq I(T(a, c), T(b, d)) \quad (46)$$

If J is an arbitrary index set and if $(a_j)_{j \in J}$ and $(b_j)_{j \in J}$ are families in $[0, 1]$, it holds that

$$T(\sup_{j \in J} a_j, b) = \sup_{j \in J} T(a_j, b) \quad (47)$$

$$I(a, \inf_{j \in J} b_j) = \inf_{j \in J} I(a, b_j) \quad (48)$$

$$T(\inf_{j \in J} a_j, b) \leq \inf_{j \in J} T(a_j, b) \quad (49)$$

$$I(a, \sup_{j \in J} b_j) \geq \sup_{j \in J} I(a, b_j) \quad (50)$$

It is easy to see that for an arbitrary t-norm T it holds that

$$I(1, b) = b \quad (51)$$

Lemma 1: Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $\beta \geq 0$; it holds that

$$L_{(\alpha_1+\alpha_2, \beta)}^{\ll}(a, b) = L_{(\alpha_1, \beta)}^{\ll}(a + \alpha_2, b) \quad (52)$$

$$L_{(\alpha_1+\alpha_2, \beta)}^{\ll}(a, b) = L_{(\alpha_1, \beta)}^{\ll}(a, b - \alpha_2) \quad (53)$$

$$L_{(\alpha_1+\alpha_2, \beta)}^{\gg}(a, b) = L_{(\alpha_1, \beta)}^{\gg}(a - \alpha_2, b) \quad (54)$$

$$L_{(\alpha_1+\alpha_2, \beta)}^{\gg}(a, b) = L_{(\alpha_1, \beta)}^{\gg}(a, b + \alpha_2) \quad (55)$$

for all a and b in \mathbb{R} .

Lemma 2: Let $\alpha \in \mathbb{R}$ and $\beta_1, \beta_2 \geq 0$; it holds that

$$L_{(\alpha+\beta_2, \beta_1)}^{\ll}(a, b) \leq L_{(\alpha+\min(\beta_1, \beta_2), \max(\beta_1, \beta_2))}^{\ll}(a, b) \quad (56)$$

for all a and b in \mathbb{R} .

Proof: If $\beta_1 \geq \beta_2$ then the proof is trivial, therefore assume that $\beta_1 < \beta_2$. For $b - a \leq \alpha + \beta_2$ and for $b - a > \alpha + \beta_1 + \beta_2$ (56) holds trivially, since in the former case the left-hand side of (56) equals 0, while in the latter case the right-hand side equals 1. Hence we only need to consider the

case where $\beta_1 > 0$ and $\alpha + \beta_2 < b - a \leq \alpha + \beta_1 + \beta_2$; it holds that

$$\begin{aligned} \frac{b - a - \alpha - \beta_2}{\beta_1} &\leq \frac{b - a - \alpha - \beta_1}{\beta_2} \\ \Leftrightarrow (b - a - \alpha)\beta_2 - \beta_2^2 &\leq (b - a - \alpha)\beta_1 - \beta_1^2 \\ \Leftrightarrow (b - a - \alpha)(\beta_2 - \beta_1) &\leq \beta_2^2 - \beta_1^2 \end{aligned}$$

Since $\beta_2 - \beta_1 > 0$ we obtain

$$\Leftrightarrow b - a \leq \alpha + \beta_1 + \beta_2$$

which completes the proof. \blacksquare

Lemma 3 (Transitivity): Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $\beta_1, \beta_2 \geq 0$; it holds that

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)) \\ \leq L_{(\alpha_1 + \alpha_2 + \min(\beta_1, \beta_2), \max(\beta_1, \beta_2))}^{\llcorner}(a, c) \end{aligned} \quad (57)$$

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\lrcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\lrcorner}(b, c)) \\ \leq L_{(\alpha_1 + \alpha_2, \max(\beta_1, \beta_2))}^{\lrcorner}(a, c) \end{aligned} \quad (58)$$

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\lrcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)) \\ \leq L_{(\alpha_2 - \alpha_1 + \min(\beta_1, \beta_2) - \beta_1, \max(\beta_1, \beta_2))}^{\llcorner}(a, c) \end{aligned} \quad (59)$$

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\lrcorner}(b, c)) \\ \leq L_{(\alpha_1 - \alpha_2 + \min(\beta_1, \beta_2) - \beta_2, \max(\beta_1, \beta_2))}^{\lrcorner}(a, c) \end{aligned} \quad (60)$$

for all a, b and c in \mathbb{R} .

Proof: As an example we prove (57), the proof of (58)–(60) is analogous. When $b - a \leq \alpha_1$ or $c - b \leq \alpha_2$ (57) obviously holds since the left-hand side equals 0. When $b - a > \alpha_1 + \beta_1$ we have

$$T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)) = L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)$$

Because the first partial mappings of $L_{(\alpha_2, \beta_2)}^{\llcorner}$ are decreasing, and by using the assumption $b - a > \alpha_1 + \beta_1$ we obtain

$$L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c) \leq L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1, c)$$

and by (52)

$$L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1, c) = L_{(\alpha_1 + \alpha_2 + \beta_1, \beta_2)}^{\llcorner}(a, c)$$

and by Lemma 2

$$L_{(\alpha_1 + \alpha_2 + \beta_1, \beta_2)}^{\llcorner}(a, c) \leq L_{(\alpha_1 + \alpha_2 + \min(\beta_1, \beta_2), \max(\beta_1, \beta_2))}^{\llcorner}(a, c)$$

In the same way, we can prove (57) when $c - b > \alpha_2 + \beta_2$. Finally, assume $\alpha_1 < b - a \leq \alpha_1 + \beta_1$ and $\alpha_2 < c - b \leq \alpha_2 + \beta_2$ (hence $\beta_1 > 0$ and $\beta_2 > 0$). For $\beta_1 \leq \beta_2$ we obtain

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)) \\ = \max(0, L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b) + L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c) - 1) \\ = \max(0, \frac{b - a - \alpha_1}{\beta_1} + \frac{c - b - \alpha_2}{\beta_2} - 1) \\ = \max(0, \frac{b - a - \alpha_1 - \beta_1}{\beta_1} + \frac{c - b - \alpha_2}{\beta_2}) \end{aligned}$$

Since $b - a - \alpha_1 - \beta_1 \leq 0$ and $\beta_1 \leq \beta_2$, we have

$$\begin{aligned} &\leq \max(0, \frac{b - a - \alpha_1 - \beta_1}{\beta_2} + \frac{c - b - \alpha_2}{\beta_2}) \\ &= \max(0, \frac{c - a - \alpha_1 - \alpha_2 - \beta_1}{\beta_2}) \\ &= L_{(\alpha_1 + \alpha_2 + \beta_1, \beta_2)}^{\llcorner}(a, c) \end{aligned}$$

For $\beta_1 > \beta_2$ the proof is entirely analogous. \blacksquare

A. Proof of Proposition 1 (Composition)

We prove (7) as an example, the proof of (8)–(10) is analogous. By (57) we already have

$$\begin{aligned} (L_{(\alpha_1, \beta_1)}^{\llcorner} \circ_{T_W} L_{(\alpha_2, \beta_2)}^{\llcorner})(a, c) \\ \leq L_{(\alpha_1 + \alpha_2 + \min(\beta_1, \beta_2), \max(\beta_1, \beta_2))}^{\llcorner}(a, c) \end{aligned}$$

for arbitrary a and c in \mathbb{R} . Reversely, for $\beta_1 \leq \beta_2$ we have

$$\begin{aligned} \sup_{b \in \mathbb{R}} T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, c)) \\ \geq \sup_{\varepsilon > 0} T_W(L_{(\alpha_1, \beta_1)}^{\llcorner}(a, a + \alpha_1 + \beta_1 + \varepsilon), \\ L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1 + \varepsilon, c)) \\ = \sup_{\varepsilon > 0} T_W(1, L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1 + \varepsilon, c)) \\ = \sup_{\varepsilon > 0} L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1 + \varepsilon, c) \end{aligned}$$

Taking into account that the first partial mappings of $L_{(\alpha_2, \beta_2)}^{\llcorner}$ are decreasing and right-continuous, we obtain

$$= L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1, c)$$

and by (52)

$$L_{(\alpha_2, \beta_2)}^{\llcorner}(a + \alpha_1 + \beta_1, c) = L_{(\alpha_1 + \alpha_2 + \beta_1, \beta_2)}^{\llcorner}(a, c)$$

For $\beta_1 > \beta_2$ the proof is entirely analogous.

B. Proof of Proposition 2

For $b - a < -\alpha_1 - \beta_1$ or $a - b \leq \alpha_2$ (15) trivially holds. Hence, we can assume that $-\alpha_1 - \beta_1 \leq b - a < -\alpha_2$, and thus, using the assumptions $\beta_1 \leq \beta_2$ and $\alpha_1 \leq \alpha_2$, that $-\alpha_1 - \beta_1 \leq b - a < -\alpha_1$ and $\alpha_2 < a - b \leq \alpha_2 + \beta_2$ (hence $\beta_1 > 0$ and $\beta_2 > 0$), we obtain

$$\begin{aligned} T_W(L_{(\alpha_1, \beta_1)}^{\lrcorner}(a, b), L_{(\alpha_2, \beta_2)}^{\llcorner}(b, a)) \\ = \max(0, L_{(\alpha_1, \beta_1)}^{\lrcorner}(a, b) + L_{(\alpha_2, \beta_2)}^{\llcorner}(b, a) - 1) \\ = \max\left(0, \frac{b - a + \alpha_1 + \beta_1}{\beta_1} + \frac{a - b - \alpha_2}{\beta_2} - 1\right) \\ = \max\left(0, \frac{b - a + \alpha_1}{\beta_1} + \frac{a - b - \alpha_2}{\beta_2}\right) \\ \leq \max\left(0, \frac{b - a + \alpha_1}{\beta_2} + \frac{a - b - \alpha_2}{\beta_2}\right) \end{aligned}$$

since $b - a + \alpha_1 < 0$. Finally, we have

$$\max\left(0, \frac{b - a + \alpha_1}{\beta_2} + \frac{a - b - \alpha_2}{\beta_2}\right) = \max\left(0, \frac{\alpha_1 - \alpha_2}{\beta_2}\right)$$

which concludes the proof since $\frac{\alpha_1 - \alpha_2}{\beta_2} \leq 0$.

APPENDIX II

PROOF OF THE TRANSITIVITY TABLE FOR RELATEDNESS MEASURES

In this appendix we provide a proof for the inequalities in Table III. Throughout this appendix, let R and S be fuzzy relations in a universe U , and let A , B and C be normalised fuzzy sets in U . Recall that the inverse fuzzy relation R^{-1} of a fuzzy relation R in U is the fuzzy relation in U defined by $R^{-1}(v, u) = R(u, v)$ for all u and v in U .

Lemma 4 (Duality):

$$A \circ_T R \circ_T B = B \circ_T R^{-1} \circ_T A \quad (61)$$

$$A \triangleleft_I R \triangleright_I B = B \triangleleft_I R^{-1} \triangleright_I A \quad (62)$$

$$(A \circ_T R) \triangleright_I B = B \triangleleft_I (R^{-1} \circ_T A) \quad (63)$$

$$A \circ_T (R \triangleright_I B) = (B \triangleleft_I R^{-1}) \circ_T A \quad (64)$$

Proof: (63) and (64) follow immediately from the definitions of the relatedness measures; (61) follows from (47), while (62) follows from (45) and (48). ■

We will now prove the correctness of the inequalities in Table III. First, note that each inequality corresponding to an entry which equals 1, trivially holds. A first series of inequalities to prove is:

$$T(A \triangleleft_I R \triangleright_I B, B \circ_T S \circ_T C) \leq (A \triangleleft_I Q) \circ_T C \quad (65)$$

$$T(A \triangleleft_I R \triangleright_I B, (B \triangleleft_I S) \circ_T C) \leq (A \triangleleft_I Q) \circ_T C \quad (66)$$

$$T(A \triangleleft_I R \triangleright_I B, B \triangleleft_I (S \circ_T C)) \leq (A \triangleleft_I Q) \circ_T C \quad (67)$$

$$T(A \circ_T R \circ_T B, B \triangleleft_I S \triangleright_I C) \leq A \circ_T (Q \triangleright_I C) \quad (68)$$

$$T((A \circ_T R) \triangleright_I B, B \triangleleft_I S \triangleright_I C) \leq A \circ_T (Q \triangleright_I C) \quad (69)$$

$$T(A \circ_T (R \triangleright_I B), B \triangleleft_I S \triangleright_I C) \leq A \circ_T (Q \triangleright_I C) \quad (70)$$

where $Q = R \circ_T S$.

Proof: By using (47) we obtain

$$\begin{aligned} & T(A \triangleleft_I R \triangleright_I B, B \circ_T S \circ_T C) \\ &= T\left(\inf_{v' \in U} I(B(v')), \inf_{u \in U} I(A(u), R(u, v'))\right), \\ & \quad \sup_{w \in U} T(C(w), \sup_{v \in U} T(B(v), S(v, w))) \\ &= \sup_{v \in U} \sup_{w \in U} T\left(\inf_{v' \in U} I(B(v')), \inf_{u \in U} I(A(u), R(u, v'))\right), \\ & \quad T(C(w), T(B(v), S(v, w))) \\ &\leq \sup_{v \in U} \sup_{w \in U} T\left(I(B(v)), \inf_{u \in U} I(A(u), R(u, v))\right), \\ & \quad T(C(w), T(B(v), S(v, w))) \\ &= \sup_{v \in U} \sup_{w \in U} T\left(T(B(v), I(B(v), \inf_{u \in U} I(A(u), R(u, v))))\right), \\ & \quad T(C(w), S(v, w)) \end{aligned}$$

By (43), we have

$$\begin{aligned} &\leq \sup_{v \in U} \sup_{w \in U} T\left(\inf_{u \in U} I(A(u), R(u, v)), T(C(w), S(v, w))\right) \\ &= \sup_{v \in U} \sup_{w \in U} T\left(T\left(\inf_{u \in U} I(A(u), R(u, v)), S(v, w)\right), C(w)\right) \end{aligned}$$

and by (49) and (42)

$$\begin{aligned} &\leq \sup_{v \in U} \sup_{w \in U} T\left(\inf_{u \in U} T(I(A(u), R(u, v)), S(v, w)), C(w)\right) \\ &\leq \sup_{v \in U} \sup_{w \in U} T\left(\inf_{u \in U} I(A(u), T(R(u, v), S(v, w))), C(w)\right) \end{aligned}$$

and finally by (47) and (50)

$$\begin{aligned} &= \sup_{v \in U} T\left(\sup_{w \in U} \inf_{u \in U} I(A(u), T(R(u, v), S(v, w))), C(w)\right) \\ &\leq \sup_{w \in U} T\left(\inf_{u \in U} \sup_{v \in U} I(A(u), T(R(u, v), S(v, w))), C(w)\right) \\ &\leq \sup_{w \in U} T\left(\inf_{u \in U} I(A(u), \sup_{v \in U} T(R(u, v), S(v, w))), C(w)\right) \end{aligned}$$

Hence we have shown that (65) holds; (66) and (67) follow from (65) by using Proposition 3. Furthermore, by using duality (68) follows from (65) (using (61), (62) and (64)), (69) follows from (67) (using (62), (63) and (64)), and (70) follows from (66) (using (62) and (64)). ■

Next, we prove the following series of inequalities:

$$T((A \circ_T R) \triangleright_I B, B \circ_T S \circ_T C) \leq A \circ_T Q \circ_T C \quad (71)$$

$$T(A \circ_T (R \triangleright_I B), B \circ_T S \circ_T C) \leq A \circ_T Q \circ_T C \quad (72)$$

$$T((A \circ_T R) \triangleright_I B, (B \triangleleft_I S) \circ_T C) \leq A \circ_T Q \circ_T C \quad (73)$$

$$T(A \circ_T (R \triangleright_I B), (B \triangleleft_I S) \circ_T C) \leq A \circ_T Q \circ_T C \quad (74)$$

$$T((A \circ_T R) \triangleright_I B, B \triangleleft_I (S \circ_T C)) \leq A \circ_T Q \circ_T C \quad (75)$$

$$T(A \circ_T (R \triangleright_I B), B \triangleleft_I (S \circ_T C)) \leq A \circ_T Q \circ_T C \quad (76)$$

$$T(A \circ_T R \circ_T B, (B \triangleleft_I S) \circ_T C) \leq A \circ_T Q \circ_T C \quad (77)$$

$$T(A \circ_T R \circ_T B, B \triangleleft_I (S \circ_T C)) \leq A \circ_T Q \circ_T C \quad (78)$$

where $Q = R \circ_T S$.

Proof: The proof of (71) is analogous to the proof of (65); (72)-(76) follow from (71) by using Proposition 3. Furthermore, by using duality (77) follows from (72) (using (61) and (64)) and (78) follows from (71) (using (61) and (63)). ■

A third series of inequalities to prove is:

$$T(A \triangleleft_I R \triangleright_I B, B \triangleleft_I S \triangleright_I C) \leq A \triangleleft_I Q \triangleright_I C \quad (79)$$

$$T((A \triangleleft_I R) \circ_T B, B \triangleleft_I S \triangleright_I C) \leq A \triangleleft_I Q \triangleright_I C \quad (80)$$

$$T(A \triangleleft_I (R \circ_T B), B \triangleleft_I S \triangleright_I C) \leq A \triangleleft_I Q \triangleright_I C \quad (81)$$

$$T(A \triangleleft_I R \triangleright_I B, (B \circ_T S) \triangleright_I C) \leq A \triangleleft_I Q \triangleright_I C \quad (82)$$

$$T(A \triangleleft_I R \triangleright_I B, B \circ_T (S \triangleright_I C)) \leq A \triangleleft_I Q \triangleright_I C \quad (83)$$

where $Q = R \circ_T S$.

Proof: The proof of (81) is analogous to the proof of (65); (79) and (80) follow from (81) by using Proposition 3. Furthermore, by using duality (82) follows from (81) (using (62) and (63)) and (83) follows from (80) (using (62) and (64)). ■

Next, we prove the following series of inequalities:

$$T((A \triangleleft_I R) \circ_T B, (B \triangleleft_I S) \circ_T C) \leq (A \triangleleft_I Q) \circ_T C \quad (84)$$

$$T(A \triangleleft_I (R \circ_T B), (B \triangleleft_I S) \circ_T C) \leq (A \triangleleft_I Q) \circ_T C \quad (85)$$

$$T((A \triangleleft_I R) \circ_T B, B \triangleleft_I (S \circ_T C)) \leq (A \triangleleft_I Q) \circ_T C \quad (86)$$

$$T(A \circ_T (R \triangleright_I B), B \circ_T (S \triangleright_I C)) \leq A \circ_T (Q \triangleright_I C) \quad (87)$$

$$T((A \circ_T R) \triangleright_I B, B \circ_T (S \triangleright_I C)) \leq A \circ_T (Q \triangleright_I C) \quad (88)$$

$$T(A \circ_T (R \triangleright_I B), (B \circ_T S) \triangleright_I C) \leq A \circ_T (Q \triangleright_I C) \quad (89)$$

where $Q = R \circ_T S$.

Proof: The proof of (85) is analogous to the proof of (65); the proof of (86) is entirely analogous. Furthermore (84) follows from (85) by using Proposition 3. Finally, by duality (87) follows from (84) (using (64)), (88) follows from (86) (using (63) and (64)), and (89) follows from (85) (using (63) and (64)). ■

Finally, we still need to prove the following inequalities:

$$T(A \triangleleft_I (R \circ_T B), B \triangleleft_I (S \circ_T C)) \leq A \triangleleft_I (Q \circ_T C) \quad (90)$$

$$T((A \circ_T R) \triangleright_I B, (B \circ_T S) \triangleright_I C) \leq (A \circ_T Q) \triangleright_I C \quad (91)$$

where $Q = R \circ_T S$.

Proof: By (49) we have

$$\begin{aligned} & T(A \triangleleft_I (R \circ_T B), B \triangleleft_I (S \circ_T C)) \\ &= T\left(\inf_{u \in U} I(A(u), \sup_{v \in U} T(B(v), R(u, v))), \right. \\ &\quad \left. \inf_{v \in U} I(B(v), \sup_{w \in U} T(C(w), S(v, w)))\right) \\ &\leq \inf_{u \in U} T\left(I(A(u), \sup_{v \in U} T(B(v), R(u, v))), \right. \\ &\quad \left. \inf_{v \in U} I(B(v), \sup_{w \in U} T(C(w), S(v, w)))\right) \end{aligned}$$

By (42) and (47) we obtain

$$\begin{aligned} &\leq \inf_{u \in U} I\left(A(u), T\left(\sup_{v \in U} T(B(v), R(u, v)), \right. \right. \\ &\quad \left. \left. \inf_{v \in U} I(B(v), \sup_{w \in U} T(C(w), S(v, w)))\right)\right) \\ &= \inf_{u \in U} I\left(A(u), \sup_{v \in U} T\left(T(B(v), R(u, v)), \right. \right. \\ &\quad \left. \left. \inf_{v' \in U} I(B(v'), \sup_{w \in U} T(C(w), S(v', w)))\right)\right) \\ &\leq \inf_{u \in U} I\left(A(u), \sup_{v \in U} T\left(T(B(v), R(u, v)), \right. \right. \\ &\quad \left. \left. I(B(v), \sup_{w \in U} T(C(w), S(v, w)))\right)\right) \end{aligned}$$

and by (43) and (47)

$$\begin{aligned} &\leq \inf_{u \in U} I\left(A(u), \sup_{v \in U} T\left(R(u, v), \sup_{w \in U} T(C(w), S(v, w))\right)\right) \\ &= \inf_{u \in U} I\left(A(u), \sup_{w \in U} T\left(C(w), \sup_{v \in U} T\left(R(u, v), S(v, w)\right)\right)\right) \end{aligned}$$

which completes the proof of (90). By duality, (91) follows from (90) (using (63)). ■

APPENDIX III

PROOF OF PROPERTIES OF THE FUZZY TEMPORAL INTERVAL RELATIONS

In this appendix we will only use the Łukasiewicz t–norm and implicator. Hence we will omit the subscripts in \circ_{T_W} , \triangleleft_{I_W} and \triangleright_{I_W} .

Lemma 5: Let A be a fuzzy set in U , B a fuzzy set in V and R a fuzzy relation from U to V . It holds that

$$1 - A \circ (R \triangleright B) = (B \circ coR^{-1}) \triangleright A \quad (92)$$

$$1 - (A \triangleleft R) \circ B = B \triangleleft (coR^{-1} \circ A) \quad (93)$$

$$1 - A \circ R \circ B = B \triangleleft coR^{-1} \triangleright A \quad (94)$$

Proof: As an example we prove (92). We obtain

$$\begin{aligned} & 1 - A \circ (R \triangleright B) \\ &= 1 - \sup_{u \in U} T_W(A(u), \inf_{v \in V} I_W(B(v), R(u, v))) \\ &= 1 - \sup_{u \in U} \max(0, A(u) \\ &\quad + \inf_{v \in V} \min(1, 1 - B(v) + R(u, v)) - 1) \\ &= 1 - \sup_{u \in U} \max(0, A(u) + \inf_{v \in V} \min(0, -B(v) + R(u, v))) \\ &= \inf_{u \in U} \min(1, 1 - A(u) + \sup_{v \in V} \max(0, B(v) - R(u, v))) \\ &= \inf_{u \in U} \min(1, 1 - A(u) \\ &\quad + \sup_{v \in V} \max(0, B(v) + coR(u, v) - 1)) \\ &= \inf_{u \in U} I_W(A(u), \sup_{v \in V} T_W(B(v), coR(u, v))) \end{aligned}$$

Lemma 6: For all a, b and c in $[0, 1]$ it holds that

$$S_W(\min(a, b), \min(a, c)) \geq \min(a, S_W(b, c)) \quad (95)$$

Proof: We have

$$\begin{aligned} & S_W(\min(a, b), \min(a, c)) \\ &= \min(1, \min(a, b) + \min(a, c)) \\ &= \min(1, a + a, a + c, a + b, b + c) \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & \min(a, S_W(b, c)) = \min(a, \min(1, b + c)) \\ &= \min(1, a, b + c) \end{aligned}$$

which proves (95) since $\min(a + a, a + b, a + c) \geq a$. ■

Lemma 7: For all a, b and c in $[0, 1]$ it holds that

$$S_W(\min(a, b), c) = \min(S_W(a, c), S_W(b, c)) \quad (96)$$

Proof: We obtain

$$\begin{aligned} & \min(S_W(a, c), S_W(b, c)) \\ &= \min(\min(1, a + c), \min(1, b + c)) \\ &= \min(1, a + c, b + c) \\ &= \min(1, \min(a, b) + c) \\ &= S_W(\min(a, b), c) \end{aligned}$$

A. Proof of Proposition 7 (Exhaustivity)

We have

$$\begin{aligned} & S_W(s_{\gamma^2}(A, B), si_{\gamma^2}(A, B), e_{\gamma^2}(A, B)) \\ &= S_W(\min((A \circ L_{\gamma^1}^{\leq}) \triangleright B, (B \circ L_{\gamma^1}^{\leq}) \triangleright A, \\ &\quad (A \triangleleft L_{\gamma^1}^{\leq}) \circ B), \\ &\quad \min((B \circ L_{\gamma^1}^{\leq}) \triangleright A, (A \circ L_{\gamma^1}^{\leq}) \triangleright B, \\ &\quad (B \triangleleft L_{\gamma^1}^{\leq}) \circ A), \\ &\quad \min((A \circ L_{\gamma^1}^{\leq}) \triangleright B, (B \circ L_{\gamma^1}^{\leq}) \triangleright A, \\ &\quad A \triangleleft (L_{\gamma^1}^{\leq} \circ B), B \triangleleft (L_{\gamma^1}^{\leq} \circ A))) \end{aligned}$$

By twice applying (95) we obtain that

$$\begin{aligned} S_W(\min(a, b_1), \min(a, b_2), \min(a, b_3)) \\ \geq \min(a, S_W(b_1, b_2, b_3)) \end{aligned}$$

for all a, b_1, b_2 and b_3 in $[0, 1]$. Substituting

$$\begin{aligned} a &:= \min((A \circ L_{\gamma_1}^{\lessdot}) \triangleright B, (B \circ L_{\gamma_1}^{\lessdot}) \triangleright A) \\ b_1 &:= (A \triangleleft L_{\gamma_1}^{\lessdot}) \circ B \\ b_2 &:= (B \triangleleft L_{\gamma_1}^{\lessdot}) \circ A \\ b_3 &:= \min(A \triangleleft (L_{\gamma_1}^{\lessdot} \circ B), B \triangleleft (L_{\gamma_1}^{\lessdot} \circ A)) \end{aligned}$$

we obtain

$$\begin{aligned} S_W(s_{\gamma^2}(A, B), si_{\gamma^2}(A, B), e_{\gamma^2}(A, B)) \\ \geq \min((A \circ L_{\gamma_1}^{\lessdot}) \triangleright B, (B \circ L_{\gamma_1}^{\lessdot}) \triangleright A, \\ S_W((A \triangleleft L_{\gamma_1}^{\lessdot}) \circ B, (B \triangleleft L_{\gamma_1}^{\lessdot}) \circ A, \\ \min(A \triangleleft (L_{\gamma_1}^{\lessdot} \circ B), \\ B \triangleleft (L_{\gamma_1}^{\lessdot} \circ A)))) \end{aligned}$$

and by (96)

$$\begin{aligned} &= \min((A \circ L_{\gamma_1}^{\lessdot}) \triangleright B, (B \circ L_{\gamma_1}^{\lessdot}) \triangleright A, \\ &S_W((A \triangleleft L_{\gamma_1}^{\lessdot}) \circ B, (B \triangleleft L_{\gamma_1}^{\lessdot}) \circ A, A \triangleleft (L_{\gamma_1}^{\lessdot} \circ B)), \\ &S_W((A \triangleleft L_{\gamma_1}^{\lessdot}) \circ B, (B \triangleleft L_{\gamma_1}^{\lessdot}) \circ A, B \triangleleft (L_{\gamma_1}^{\lessdot} \circ A))) \end{aligned}$$

and by (93) and (2)

$$\begin{aligned} &= \min((A \circ L_{\gamma_1}^{\lessdot}) \triangleright B, (B \circ L_{\gamma_1}^{\lessdot}) \triangleright A) \\ &= hh_{\gamma^1}(A, B) \end{aligned}$$

since $S_W(a, 1 - a) = 1$ for all a in $[0, 1]$. We can show analogously that

$$\begin{aligned} S_W(o_{\gamma^3}(A, B), fi_{\gamma^2}(A, B), di_{\gamma^2}(A, B)) \\ \geq \min(B \circ (L_{\gamma_1}^{\lessdot} \triangleright A), A \circ L_{\gamma_1}^{\lessdot} \circ B) \\ = oc_{\gamma^2}(A, B) \end{aligned}$$

$$\begin{aligned} S_W(oi_{\gamma^3}(A, B), f_{\gamma^2}(A, B), d_{\gamma^2}(A, B)) \\ \geq \min(A \circ (L_{\gamma_1}^{\lessdot} \triangleright B), B \circ L_{\gamma_1}^{\lessdot} \circ A) \\ = yc_{\gamma^2}(A, B) \end{aligned}$$

$$\begin{aligned} S_W(b_{\gamma^1}(A, B), m_{\gamma^1}(A, B)) \\ \geq A \triangleleft L_{\gamma_1}^{\lessdot} \triangleright B \\ = pr_{\gamma^1}(A, B) \end{aligned}$$

$$\begin{aligned} S_W(bi_{\gamma^1}(A, B), mi_{\gamma^1}(A, B)) \\ \geq B \triangleleft L_{\gamma_1}^{\lessdot} \triangleright A \\ = sd_{\gamma^1}(A, B) \end{aligned}$$

$$\begin{aligned} S_W(hh_{\gamma^1}(A, B), yc_{\gamma^2}(A, B), oc_{\gamma^2}(A, B)) \\ \geq \min(A \circ L_{\gamma_1}^{\lessdot} \circ B, B \circ L_{\gamma_1}^{\lessdot} \circ A) \\ = ct_{\gamma^2}(A, B) \end{aligned}$$

$$S_W(ct_{\gamma^2}(A, B), pr_{\gamma^1}(A, B), sd_{\gamma^1}(A, B)) = 1$$

which completes the proof. Note that the conceptual neighborhoods pr and sd are not included in Table VI as they are not relevant to the transitivity table for temporal interval relations. Here the (generalized) conceptual neighborhoods are merely used as a shorthand.

B. Proof of Proposition 8 (Mutual Exclusiveness)

To prove the mutual exclusiveness of the fuzzy temporal relations, 78 cases have to be considered. Here, as an example, we provide a proof for two of these cases. First, we show that

$$T_W(d_{(\alpha, \beta, \alpha, \beta)}(A, B), m_{(\alpha, \beta)}(A, B)) = 0$$

Using Table III we obtain

$$\begin{aligned} T_W(d_{(\alpha, \beta, \alpha, \beta)}(A, B), m_{(\alpha, \beta)}(A, B)) \\ = T_W(\min(B \circ (L_{\gamma}^{\lessdot} \triangleright A), (A \triangleleft L_{\gamma}^{\lessdot}) \circ B), \\ \min(A \triangleleft L_{\gamma}^{\lessdot} \triangleright B, B \circ L_{\gamma}^{\lessdot} \circ A)) \\ \leq T_W(B \circ (L_{\gamma}^{\lessdot} \triangleright A), A \triangleleft L_{\gamma}^{\lessdot} \triangleright B) \\ \leq B \circ ((L_{\gamma}^{\lessdot} \circ L_{\gamma}^{\lessdot}) \triangleright B) \end{aligned}$$

where $\gamma = (\alpha, \beta)$. By (10) we obtain

$$= B \circ (L_{(0, \beta)}^{\lessdot} \triangleright B)$$

which equals 0 by Proposition 5, since for $\alpha \geq 0$, $L_{(\alpha, \beta)}^{\lessdot}$ is an irreflexive fuzzy relation.

As a second example, we show that

$$T_W(s_{\gamma^2}(A, B), m_{\gamma^1}(A, B)) = 0$$

Since A is nondegenerate w.r.t. $(2\alpha, \beta)$ we obtain using Table III and (9)

$$\begin{aligned} (B \circ L_{\gamma}^{\lessdot}) \triangleright A \\ = T_W((B \circ L_{\gamma}^{\lessdot}) \triangleright A, A \circ L_{(2\alpha, \beta)}^{\lessdot} \circ A) \\ \leq B \circ (L_{\gamma}^{\lessdot} \circ L_{(2\alpha, \beta)}^{\lessdot}) \circ A \\ = B \circ L_{(\alpha, \beta)}^{\lessdot} \circ A \end{aligned}$$

We obtain using Table III and (10)

$$\begin{aligned} T_W(s_{\gamma^2}(A, B), m_{\gamma^1}(A, B)) \\ = T_W(\min((A \circ L_{\gamma}^{\lessdot}) \triangleright B, (B \circ L_{\gamma}^{\lessdot}) \triangleright A, \\ (A \triangleleft L_{\gamma}^{\lessdot}) \circ B), \\ \min(A \triangleleft L_{\gamma}^{\lessdot} \triangleright B, B \circ L_{\gamma}^{\lessdot} \circ A)) \\ \leq T_W((B \circ L_{\gamma}^{\lessdot}) \triangleright A, A \triangleleft L_{\gamma}^{\lessdot} \triangleright B) \\ \leq T_W(B \circ L_{\gamma}^{\lessdot} \circ A, A \triangleleft L_{\gamma}^{\lessdot} \triangleright B) \\ \leq B \circ ((L_{\gamma}^{\lessdot} \circ L_{\gamma}^{\lessdot}) \triangleright B) \\ = B \circ (L_{(0, \beta)}^{\lessdot} \triangleright B) \end{aligned}$$

which equals 0 by Proposition 5, since for $\alpha \geq 0$, $L_{(\alpha, \beta)}^{\lessdot}$ is an irreflexive fuzzy relation.

C. Proof of Proposition 9 ((A)symmetry)

As an example, we show that

$$T_W(d_{\gamma_{12}}(A, B), d_{\gamma_{12}}(B, A)) = 0$$

By using Table III and (7) we obtain

$$\begin{aligned} T_W(d_{\gamma_{12}}(A, B), d_{\gamma_{12}}(B, A)) &= T_W(\min((A \triangleleft L_{\gamma_2}^{\ll} \circ B, B \circ (L_{\gamma_1}^{\ll} \triangleright A)), \\ &\quad \min((B \triangleleft L_{\gamma_2}^{\ll} \circ A, A \circ (L_{\gamma_1}^{\ll} \triangleright B)))) \\ &\leq T_W((A \triangleleft L_{\gamma_2}^{\ll} \circ B, (B \triangleleft L_{\gamma_2}^{\ll} \circ A)) \\ &\leq (A \triangleleft (L_{\gamma_2}^{\ll} \circ L_{\gamma_2}^{\ll})) \circ A \\ &= (A \triangleleft L_{(2\alpha_2 + \beta_2, \beta_2)}^{\ll}) \circ A \end{aligned}$$

which equals 0 by Proposition 5, since for $\alpha \geq 0$, $L_{(\alpha, \beta)}^{\ll}$ is an irreflexive fuzzy relation.

As another example, we show that

$$T_W(m_{(\alpha_1, \beta_1)}(A, B), m_{(\alpha_1, \beta_1)}(B, A)) = 0$$

if A and B are nondegenerate fuzzy time periods w.r.t. (α_1, β_1) . We obtain

$$\begin{aligned} T_W(m_{(\alpha_1, \beta_1)}(A, B), m_{(\alpha_1, \beta_1)}(B, A)) &= T_W(\min(A \triangleleft L_{\gamma_1}^{\ll} \triangleright B, B \circ L_{\gamma_1}^{\ll} \circ A), \\ &\quad \min(B \triangleleft L_{\gamma_1}^{\ll} \triangleright A, A \circ L_{\gamma_1}^{\ll} \circ B)) \\ &\leq T_W(A \triangleleft L_{\gamma_1}^{\ll} \triangleright B, B \triangleleft L_{\gamma_1}^{\ll} \triangleright A) \end{aligned}$$

Since A is nondegenerate, we have by Table III and (10) that

$$\begin{aligned} A \triangleleft L_{\gamma_1}^{\ll} \triangleright B &= T_W(A \circ L_{\gamma_1}^{\ll} \circ A, A \triangleleft L_{\gamma_1}^{\ll} \triangleright B) \\ &\leq A \circ ((L_{\gamma_1}^{\ll} \circ L_{\gamma_1}^{\ll}) \triangleright B) \\ &= A \circ (L_{(0, \beta_1)}^{\ll} \triangleright B) \end{aligned}$$

Analogously, since B is nondegenerate, we obtain

$$B \triangleleft L_{\gamma_1}^{\ll} \triangleright A \leq B \circ (L_{(0, \beta_1)}^{\ll} \triangleright A)$$

Hence we already have

$$\begin{aligned} T_W(m_{(\alpha_1, \beta_1)}(A, B), m_{(\alpha_1, \beta_1)}(B, A)) &\leq T_W(A \circ (L_{(0, \beta_1)}^{\ll} \triangleright B), B \circ (L_{(0, \beta_1)}^{\ll} \triangleright A)) \end{aligned}$$

and by Table III and (7)

$$\begin{aligned} &\leq A \circ ((L_{(0, \beta_1)}^{\ll} \circ L_{(0, \beta_1)}^{\ll}) \triangleright A) \\ &= A \circ (L_{(\beta_1, \beta_1)}^{\ll} \triangleright A) \end{aligned}$$

which equals 0 by Proposition 5, since for $\alpha \geq 0$, $L_{(\alpha, \beta)}^{\ll}$ is an irreflexive fuzzy relation.

D. Proof of Proposition 10 ((Ir)reflexivity)

We will only prove that $m_{\gamma_1}(A, A) = 0$ when A is a nondegenerate fuzzy time period w.r.t. (α_1, β_1) , as the other equalities follow straightforwardly from Proposition 4 and Proposition 5. We obtain

$$\begin{aligned} m_{\gamma_1}(A, A) &= \min(A \triangleleft L_{\gamma_1}^{\ll} \triangleright A, A \circ L_{\gamma_1}^{\ll} \circ A) \\ &\leq A \triangleleft L_{\gamma_1}^{\ll} \triangleright A \end{aligned}$$

Since A is nondegenerate w.r.t. (α_1, β_1) we have by Table III and (9)

$$\begin{aligned} &= T_W(A \triangleleft L_{\gamma_1}^{\ll} \triangleright A, A \circ L_{\gamma_1}^{\ll} \circ A) \\ &\leq (A \triangleleft (L_{\gamma_1}^{\ll} \circ L_{\gamma_1}^{\ll})) \circ A \\ &= (A \triangleleft L_{(0, \beta_1)}^{\ll}) \circ A \end{aligned}$$

which equals 0 by Proposition 5.

ACKNOWLEDGMENT

Steven Schockaert would like to thank the Research Foundation – Flanders for funding his research.

REFERENCES

- [1] D. Ahn, S. Fissaha Adafre, and M. de Rijke, "Extracting temporal information from open domain text: a comparative exploration," *J. Digital Information Management*, vol. 3, no. 1, pp. 14–20, 2005.
- [2] D. Ahn, S. Schockaert, M. De Cock, and E.E. Kerre, "Supporting temporal question answering: strategies for offline data collection," *Proc. of the 5th conf. on Inference in Computational Semantics*, available at <http://www.cs.man.ac.uk/~ipratt/ICOS-5/ICOS-5-PDF/Ahn.pdf>, 2006.
- [3] J.F. Allen, "Maintaining knowledge about temporal intervals," *Communications of the ACM*, vol. 26, no. 11, pp. 832–843, 1983.
- [4] S. Badaloni and M. Giacomini, "A fuzzy extension of Allen's interval algebra," *Lecture Notes in Artificial Intelligence*, vol. 1792, pp. 155–165, 2000.
- [5] W. Bandler and L.J. Kohout, "Fuzzy relational products as a tool for analysis and synthesis of the behaviour of complex natural and artificial systems," *Fuzzy Sets: Theory and Application to Policy Analysis and Information Systems (S.K. Wang and P.P. Chang, eds.)*, Plenum Press, New York and London, pp. 341–367, 1980.
- [6] B.D. Bašić and S. Ribarić, "A new approach for modelling fuzzy time primitives," *Proc. of the 10th Mediterranean Electrotechnical Conf.*, vol. 2, pp. 721–724, 2000.
- [7] T. Bittner, "Approximate qualitative temporal reasoning," *Annals of Mathematics and Artificial Intelligence*, vol. 35, no. 1–2, pp. 39–80, 2002.
- [8] U. Bodenhofer, "A new approach to fuzzy orderings," *Tatra Mountains Mathematical Publications*, vol. 16, pp. 21–29, 1999.
- [9] D. Dubois and H. Prade, "Processing fuzzy temporal knowledge," *IEEE Transactions on Systems, Man, and Cybernetics*, vol. 19, no. 4, pp. 729–744, 1989.
- [10] D. Dubois, A. HadjAli, and H. Prade, "Fuzziness and uncertainty in temporal reasoning," *Journal of Universal Computer Science*, vol. 9, pp. 1168–1194, 2003.
- [11] S. Dutta, "An event-based fuzzy temporal logic," *Proc. of the 18th IEEE Int. Symp. on Multiple-Valued Logic*, pp. 64–71, 1988.
- [12] E. Filatova and E. Hovy, "Assigning time-stamps to event-clauses," *Proc. of the 2001 ACL Workshop on Temporal and Spatial Information Processing*, pp. 88–95, 2001.
- [13] C. Freksa, "Temporal Reasoning Based on Semi-Intervals," *Artificial Intelligence*, vol. 54, no. 1–2, pp. 199–227, 1992.
- [14] H.W. Guesgen, J. Hertzberg, and A. Philpott, "Towards implementing fuzzy Allen relations," *Proc. of the ECAI-94 Workshop on Spatial and Temporal Reasoning*, pp. 49–55, 1994.
- [15] M. Lapata and A. Lascarides, "Inferring sentence-internal temporal relations," *Proc. of the North American Chapter of the ACL*, pp. 153–160, 2004.
- [16] I. Mani, B. Schiffman, and J. Zhang, "Inferring temporal ordering of events in news," *Proc. of the 2003 Human Language Technology Conference of the North American Chapter of the ACL*, pp. 55–57, 2003.
- [17] G. Nagypál and B. Motik, "A fuzzy model for representing uncertain, subjective and vague temporal knowledge in ontologies," *Lecture Notes in Computer Science*, vol. 2888, pp. 906–923, 2003.
- [18] B. Nebel and H.-J. Bürckert, "Reasoning about temporal relations: a maximal tractable subclass of Allen's interval algebra," *Journal of the ACM*, vol. 42, no. 1, pp. 43–66, 1995.
- [19] V. Novák, I. Perfilieva, and J. Močkoř, *Mathematical principles of fuzzy logic*, Kluwer Academic Publishers, 1999.

- [20] H.J. Ohlbach, "Relations between fuzzy time intervals," *Proc. of the 11th Int. Symp. on Temporal Representation and Reasoning*, pp. 44–51, 2004.
- [21] D.A. Randell, Z. Cui, and A.G. Cohn, "A spatial logic based on regions and connection," *Proc. of the 3rd Int. Conf. on Principles of Knowledge Representation and Reasoning*, pp. 165–176, 1992.
- [22] E. Saquete, P. Martínez-Barco, R. Muñoz, and J.L. Vicedo, "Splitting complex temporal questions for question answering systems," *Proc. of the 42nd Annual Meeting of the ACL*, pp. 566–573, 2004.
- [23] F. Schilder and C. Habel, "From temporal expressions to temporal information: semantic tagging of news messages," *Proc. of 2001 ACL Workshop for Temporal and Spatial Information Processing*, pp. 65–72, 2001.
- [24] F. Schilder and C. Habel, "Temporal information extraction for temporal question answering," *Proc. of the 2003 AAAI Spring Symposium*, pp. 35–44, 2003.
- [25] F. Schilder, "Extracting meaning from temporal nouns and temporal prepositions," *ACM Transactions on Asian Language Information Processing*, vol. 3, no. 1, pp. 33–50, 2004.
- [26] S. Schockaert, D. Ahn, M. De Cock, and E.E. Kerre, "Question answering with imperfect temporal information," *Proc. 7th Int. Conf. on Flexible Query Answering Systems*, LNAI 4027, pp. 647–658, 2006.
- [27] S. Schockaert, "Construction of membership functions for fuzzy time periods," *Proc. of the ESSLLI 2005 Student Session*, pp. 305–317, 2005.
- [28] S. Schockaert, M. De Cock, and C. Cornelis, "Relatedness of fuzzy sets," *Journal of Intelligent and Fuzzy Systems*, vol. 6, no. 4, pp. 297–303, 2005.
- [29] S. Schockaert, M. De Cock, and E.E. Kerre, "Imprecise temporal interval relations," *Lecture Notes in Computer Science*, vol. 3849, pp. 108–113, 2006.
- [30] S. Schockaert, M. De Cock, and E.E. Kerre, "An Efficient Characterization of Fuzzy Temporal Interval Relations," *Proc. of the 2006 IEEE World Congress on Computational Intelligence*, pp. 9026–9033, 2006.
- [31] P. van Beek, "Reasoning about qualitative temporal information," *Artificial Intelligence*, vol. 58, pp. 297–326, 1992.
- [32] M. Vilain, H. Kautz, and P. van Beek, "Constraint propagation algorithms for temporal reasoning: a revised report," *Readings in qualitative reasoning about physical systems (D.S. Weld and J. de Kleer, eds.)*, Morgan Kaufmann, San Mateo, pp. 373–381, 1989.
- [33] L.A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no. 3, pp. 338–353, 1965.



Steven Schockaert obtained a M.Sc. degree in Computer Science at Ghent University in 2004. Since October 1, 2004, he is a research assistant supported by the Research Foundation – Flanders. His research primarily deals with intelligent techniques for focused information retrieval. In particular, the use of temporal representation and reasoning to process questions with temporal restrictions in question answering systems, and the use of spatial representation and reasoning to increase the coverage and flexibility of geographical information retrieval

systems. He is currently preparing a Ph.D. thesis under the supervision of Etienne E. Kerre and Martine De Cock.



Martine De Cock holds a M.Sc. and a Ph.D. degree in Computer Science from Ghent University. From 1998 until 2005 her activities as a research assistant and a postdoctoral fellow were supported by the Research Foundation – Flanders. She worked as a visiting scholar in the BISC group at the University of California, Berkeley (USA) and the Knowledge Systems Laboratory at Stanford University (USA). Since 2005 she is a professor at the Department of Applied Mathematics and Computer Science.

Her current research efforts are directed towards the development and the use of computational intelligent methods for next generation web applications.



Etienne E. Kerre obtained his M.Sc. degree in Mathematics in 1967 and his Ph.D. in Mathematics in 1970 from Ghent University. Since 1984, he has been a lecturer, and since 1991, a full professor at Ghent University. He is a referee for 50 international scientific journals, and also a member of the editorial board of international journals and conferences on fuzzy set theory. He was an honorary chairman at various international conferences. In 1976, he founded the Fuzziness and Uncertainty Modelling Research Unit (FUM) and since then his research

has been focused on the modelling of fuzziness and uncertainty, and has resulted in a great number of contributions in fuzzy set theory and its various generalizations. Especially the theories of fuzzy relational calculus and of fuzzy mathematical structures owe a great deal to him. Over the years he has also been a supervisor of 20 Ph.D.'s on fuzzy set theory. His current research interests include fuzzy and intuitionistic fuzzy relations, fuzzy topology, and fuzzy image processing. He has authored or co-authored 11 books, and more than 350 papers appeared in international refereed journals and proceedings.

LIST OF FIGURES

1	Fuzzy ordering of time points	3
2	Fuzzy time intervals A and B	6
3	Modelling the time span of the Cold War	11

LIST OF TABLES

I	Allen's temporal interval relations between intervals $A = [a^-, a^+]$ and $B = [b^-, b^+]$	1
II	Relation between the boundaries of the crisp intervals $[a^-, a^+]$ and $[b^-, b^+]$, and the fuzzy intervals A and B	4
III	Transitivity table for relatedness measures.	6
IV	Fuzzy temporal interval relations. $\gamma_{123} = (\alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \gamma_3)$, $\gamma_{12} = (\alpha_1, \beta_1, \alpha_2, \beta_2)$, $\gamma_1 = (\alpha_1, \beta_1)$, $\gamma_2 = (\alpha_2, \beta_2)$ and $\gamma_3 = (\alpha_3, \beta_3)$	6
V	Transitivity table for fuzzy temporal interval relations ($\gamma^1 = (0, 0)$, $\gamma^2 = (0, 0, 0, 0)$, $\gamma^3 = (0, 0, 0, 0, 0, 0)$) . Let $\mathcal{R}_{(R_1, R_2)}$ be the entry in this table on the row corresponding with the fuzzy temporal relation R_1 and the column corresponding with the fuzzy temporal relation R_2 . For nondegenerate (w.r.t. $(0, 0)$) fuzzy time periods A , B and C , it holds that $T_W(R_1(A, B), R_2(B, C)) \leq \mathcal{R}_{(R_1, R_2)}(A, C)$	9
VI	Freksa's conceptual neighborhoods [13].	10